

# in Neuroscience (<https://www.frontiersin.org/journals/in neuroscience>)

THIS ARTICLE IS PART OF THE RESEARCH TOPIC  
Nanotechnologies in Neuroscience and Neuroengineering [View all 29 Articles >](#)  
(<https://www.frontiersin.org/research-topics/3361#articles>)

## TABLE OF CONTENTS

- Abstract
- Introduction
- Exosomes' as a Biological Communication Tool
- Neurovascular Unit Remodeling in Response to Stroke
- Neurovascular Unit-Derived Exosomes in Response to Stroke
- The Potential Use of Mesenchymal Stem Cells and Their Exosomes for Stroke Therapy
- Conclusion and Perspectives
- Author Contributions
- Funding
- Conflict of Interest Statement



# frontiers SPOTLIGHT

([https://spotlight.frontiersin.org/how-to-enter/?utm\\_source=FWEB&utm\\_medium=FMAIN&utm\\_campaign=SPOTLIGHT&utm\\_term=BANNER&utm\\_content=WEB](https://spotlight.frontiersin.org/how-to-enter/?utm_source=FWEB&utm_medium=FMAIN&utm_campaign=SPOTLIGHT&utm_term=BANNER&utm_content=WEB))  
References

Want to win \$100,000 to host your own conference? ([https://spotlight.frontiersin.org/how-to-enter/?utm\\_source=FWEB&utm\\_medium=FMAIN&utm\\_campaign=SPOTLIGHT&utm\\_term=BANNER&utm\\_content=WEB](https://spotlight.frontiersin.org/how-to-enter/?utm_source=FWEB&utm_medium=FMAIN&utm_campaign=SPOTLIGHT&utm_term=BANNER&utm_content=WEB))

Suggest a Research Topic ([https://spotlight.frontiersin.org/how-to-enter/?utm\\_source=FWEB&utm\\_medium=FMAIN&utm\\_campaign=SPOTLIGHT&utm\\_term=BANNER&utm\\_content=WEB](https://spotlight.frontiersin.org/how-to-enter/?utm_source=FWEB&utm_medium=FMAIN&utm_campaign=SPOTLIGHT&utm_term=BANNER&utm_content=WEB))



Download Article



Export citation



4,013  
TOTAL VIEWS

SHARE ON

0 0 0 0 New

**MINI REVIEW ARTICLE**

Front. Neurosci., 06 November 2018 | https://doi.org/10.3389/fnins.2018.00811 (https://doi.org/10.3389/fnins.2018.00811)



# Multicellular Crosstalk Between Exosomes and the Neurovascular Unit After Cerebral Ischemia. Therapeutic Implications

**TABLE OF CONTENTS****Abstract****Introduction**

Exosomes' as a

**Ana-Maria Zagrean** (<http://www.frontiersin.org/people/u/165431>)<sup>1,†</sup>, **Dirk M. Hermann** (<http://www.frontiersin.org/people/u/111334>)<sup>2,3†</sup>, **Ioan Opris** (<http://www.frontiersin.org/people/u/25451>)<sup>4</sup>, **Leon Zagrean** (<http://www.frontiersin.org/people/u/164809>)<sup>1</sup> and **Aurel Popa-Wagner** (<http://www.frontiersin.org/people/u/174569>)<sup>2,3,5\*</sup>

Remodeling in

<sup>1</sup>Division of Physiology and Neuroscience, Carol Davila University of Medicine and Pharmacy, Bucharest, Romania<sup>2</sup>Department of Neurology, Chair of Vascular Neurology, Dementia and Ageing Research, University Hospital Essen, Essen, Germany  
Neurovascular Unit<sup>3</sup>Center of Clinical and Experimental Medicine, University of Medicine and Pharmacy of Craiova, Craiova, Romania

Derived Exosomes

<sup>4</sup>Department of Neurological Surgery, University of Miami, Miami, FL, United States

in Response to

<sup>5</sup>School of Medicine, Griffith University, Gold Coast, QLD, Australia**The Potential Use of**

**Restorative Strategies** after stroke are focused on the remodeling of cerebral endothelial cells and brain parenchymal cells. The latter, i.e., neurons, neural precursor cells and glial cells, synergistically interact with endothelial cells in the ischemic brain, providing a neurovascular unit (NVU) remodeling that can be used as target for stroke therapies. Intercellular communication and signaling within the NVU, the multicellular brain–vessel–blood interface, including its highly selective blood-brain barrier, are fundamental to the central nervous system homeostasis and function. Emerging research designates cell-derived extracellular vesicles and especially the nano-sized exosomes, as a complex mean of cell-to-cell communication, with potential use for clinical applications. Through their richness in active molecules and biological information (e.g., proteins, lipids, genetic material), exosomes contribute to intercellular signaling, a condition particularly required in the central nervous system. Cerebral endothelial cells, perivascular astrocytes, pericytes, microglia and neurons, all part of the NVU, have been shown to release and uptake exosomes. Also, exosomes cross the blood-brain and blood-cerebrospinal fluid barriers, allowing communication between periphery and brain, in normal and disease conditions. As such exosomes might be a powerful diagnostic tool and a promising therapeutic shuttle of natural nanoparticles, but also a means of disease spreading (e.g., immune system modulation, pro-inflammatory action, propagation of neurodegenerative factors). This review highlights the importance of exosomes in mediating the intercellular crosstalk within the NVU and reveals the restorative therapeutic potential of exosomes harvested from multipotent mesenchymal stem cells in ischemic stroke, a frequent neurologic condition lacking an efficient therapy.

**Introduction**

At the interface with the bloodstream, neurovascular units (NVUs) are structural and functional multicellular modules consisting of neurons, perivascular astrocytes, microglia, pericytes, extracellular matrix and the endothelial cells of the brain microcirculation. They provide a coordinated neurovascular coupling and maintain a highly selective blood-brain barrier (BBB) (Abbott, 2002). The dynamic multicellular crosstalk within the NVUs in physiological and pathological conditions could reveal novel cell-targeted therapeutic strategies with impact on the BBB, cerebral homeostasis and brain functions (Attwell et al., 2010; Abbott and Friedman, 2012).

The endothelial cells of the BBB are interconnected by tight and adherens junctions and form a continuous layer. This layer selectively buffers the impact of fluctuations in blood composition on brain interstitial fluid, regulating the brain microenvironment and neuronal signaling (Abbott, 2013). Various transcellular transport systems across the BBB have been described, as carrier mediated transport, receptor-mediated transport, ion transfer, efflux carriage, adsorptive-mediated passage, and fluid-phase endocytosis (Zlokovic, 2008).

Apart from the classical modes of intercellular communication, such as ligand-receptor interactions, direct cell-cell contacts (e.g., gap junctions) or paracrine signaling (Goodenough et al., 1996), a significant experimental evidence has confirmed that several physiological and pathophysiological processes are controlled by the extracellular membrane vesicles, such as exosomes and microvesicles, secreted from various cellular sources into the body fluids and interconnecting cells without direct cell-to-cell contact (Valadi et al., 2007). This type of signaling occurs mainly through exosomes, which are nano-sized vesicles that easily transfer biological information from cell to cell. This is achieved by means of exosomal molecules that would usually not cross membrane barriers. This shows the capability of inducing functional changes in target cells and modulating local and systemic crosstalk (Krämer-Albers and Hill, 2016). In the brain, exosomes are released from all types of cells (Fröhbeis et al., 2013) and are bidirectionally transported through the blood-brain communication interfaces, blood-brain and blood-cerebrospinal fluid barriers (Balusu et al., 2016). These blood-brain interfaces are potential pathways for therapeutically administered exosomes.

**Response to Stroke**  
Given their capacity to easily reach body compartments and connect origin cells with target cells, exosomes have a promising potential to be used in clinical applications. Indeed, exosomes have shown the capacity to serve both as biomarkers and novel therapeutic tools in the nervous system pathologies lacking efficient therapies, such as stroke (Bokde and Vassalli, 2017). The cellular interactions within NVU might contribute to (i) the restoration of a well-organized cerebral microvasculature by providing trophic support and a stimulating brain microenvironment (Hermann and Chopp, 2012); and (ii) the remodeling of parenchymal tissue, including axonal sprouting, dendritic growth and synaptic reorganization (Hermann and Chopp, 2012). However, there is still more to explore about the diagnostic benefits and therapeutic roles of exosomes, their production, release, transport, uptake, signaling potential, change of their cargo proteins profile and miRNAs (Zhang and Chopp, 2016). Here, we review the roles of exosomes in mediating the intercellular crosstalk within the NVU and the therapeutic potential of exosomes derived from multipotent mesenchymal stem cells (MSCs) in stroke.

**Contributions**

## Funding

## Exosomes' as a Biological Communication Tool

**Conflict of Interest**

Exosomes are defined as 30–100 nm sized membrane vesicles derivatives of the endosomal compartment and correspond to the intraluminal vesicles of multivesicular bodies (MVBs) that upon fusion of the MVBs with the plasma membrane are released as exosomes into the extracellular environment (Lener et al., 2015), where they act as signaling organelles for intercellular communication. From the extracellular milieu, exosomes may contact target cells by (i) receptor-mediated adhesion to the cellular plasma membrane, followed by endocytic uptake and internalization, (ii) direct fusion of the exosome membrane with the target cell membrane and subsequent exosomal content release into the recipient cell (Bang and Thum, 2012).

Exosomes' vesicles are homogenous in shape, surrounded by a phospholipid membrane displaying membrane proteins, such as cell-specific receptors, and containing cell-type specific combinations of lipids, metabolites, coding and non-coding RNAs (miRNA, sRNA), single- and double stranded DNA, cytosolic and membrane proteins including enzymes, growth factors, receptors and cytokines (Théry et al., 2001; Lener et al., 2015). Exosomal lipids (e.g., phosphoglycerides, sphingomyelin, cholesterol, ceramide) are important for providing structural stability. Proteins of the exosomes are characteristic for their endosomal origin, and include membrane transport and fusion proteins (annexins, flotillin), proteins involved in cell targeting (tetraspanins, mostly CD9 and CD63) or other proteins correlated with their

biogenesis from MVBS, as the tumor susceptibility gene 101 (TSG101) (András and Toborek, 2016). Exosomes also contain heat-shock proteins (Hsp60, Hsp70, Hsp90), known for their neuroprotective potential. Also, they expose low levels of phosphatidylserine and cell-type-specific proteins. One of the most important function of the exosomes is targeting cellular pathways in the recipient cells through their RNAs and miRNAs cargo (Ling et al., 2013).

Novel research supports exosomes as a fundamental mechanism of communication in the nervous system, with roles in brain homeostasis and plasticity (Holm et al., 2018), acting as bidirectional cargo in brain-periphery communication and within the brain, in between neurons, glia, vascular and perivascular cells (Figure 1). Exosome secretion has been described from (i) depolarized/stimulated cortical neurons, mainly from the somato-dendritic compartments (Faure et al., 2006; Lachenal et al., 2011; Von Bartheld and Altick, 2011), (ii) oligodendrocytes (Frühbeis et al., 2013), (iii) microglia (Potolicchio et al., 2005), (iv) astrocytes when activated by oxidative and heat stress (Taylor et al., 2007), (v) endothelial cells (Dozio and Sanchez, 2017), and (vi) pericytes (Mayo and Bearden, 2015), known to generate MSCs in the perivascular area of the lesioned or inflamed vessels (Caplan, 2008; Caplan, 2016).

## TABLE OF CONTENTS

### FIGURE 1

#### Abstract



([https://www.frontiersin.org/files/Articles/411939/fnins-12-00811-HTML/image\\_m/fnins-12-00811-g001.jpg](https://www.frontiersin.org/files/Articles/411939/fnins-12-00811-HTML/image_m/fnins-12-00811-g001.jpg))

**FIGURE 1.** Exosomes as a fundamental mechanism of communication between components of the neurovascular unit.

#### Tool

#### Neurovascular Unit

#### Remodeling in

#### Response to Stroke

The complex and versatile exosomal signaling was shown to impact the synaptic activity (e.g., neuronal origin exosomes Neurovascular Unit, exhibiting neurotransmitter receptors bind neurotransmitters within the synapse to stop signaling), *trans-synaptic communication*, synaptic plasticity, maintenance of myelination, angiogenesis, neurovascular integrity, but also on neuroregeneration and neuroprotection in response to disease conditions (Holm et al., 2018). For example, angiogenesis could be stimulated both by activation of signaling pathways PI3K, ERK1/2, Wnt4/β-catenin or NF-κB and transfer of Mesenchymal Stem Cells and Their transcription factors STAT3, STAT5, transfer of lipids like S1P, transfer of proteins including VEGF, FGF-2, PDGF, metalloproteases, but also by the transfer of micro-RNA-126, miR-214, miR-296, and miR-150 (for a review, see Todorova et al., 2017). Likewise, after experimental stroke, treatment with exosomes isolated from miR-133b-overexpressing MSCs, significantly increased functional improvement and neurite remodeling/brain plasticity in the ischemic boundary area compared with control animals (Xin et al., 2017b). Recently, it was also reported that miR-26a is a physiological regulator of mammalian axon regeneration by targeting glycogen synthase kinase 3β (GSK3β) in adult mouse sensory neurons *in vitro* and *in vivo* (Tsenkina et al., 2015).

#### Funding

Exosomes can also propagate inflammation across the BBB and within the brain, as brain endothelial cells activated by systemic inflammation further activate the neighboring cells in the NVU via secreted exosomes (Balusu et al., 2016; Holm et al., 2018). The pathogenic role of microglia-derived exosomes in the inflammatory response was demonstrated in a model of traumatic brain injury (TBI) i.e., *in vitro* activated microglia-derived exosomes induced

neuroinflammation at the site of injection and around the lesion. Furthermore, circulating enriched exosomes or CD11b-isolated microglia from the TBI brain *ex vivo*, initiated neuroinflammation following intracortical injection in naïve animals (Verderio et al., 2012; Kumar et al., 2017). The pathogenic effects of microglia-derived exosomes could be mediated by pro-inflammatory mediators TNF-alpha, IL-1β and miR-155 (Kumar et al., 2017).

Also, exosomes contribute to disease spreading by acting like Trojan horses for neurodegenerative agents (e.g., toxins, such as tetanus toxin, protein aggregates, such as phosphorylated Tau, amyloid Aβ or synuclein) (Bellingham et al., 2012; Holm et al., 2018). Through their non-coding RNA cargo and miRNA transfer, exosomes are involved in epigenetic regulation of neuro-glial communication within the nervous system, but also in brain-body epigenetic interconnection (Lai and Breakefield, 2012).

## Neurovascular Unit Remodeling in Response to Stroke

We use cookies on this website to improve your user experience. [Learn more \(//www.frontiersin.org/legal/list-of-cookies\)](https://www.frontiersin.org/legal/list-of-cookies)

CLOSE

Following the failure of acute neuroprotection therapies, major efforts are currently made worldwide to promote neurological recovery and brain plasticity in the subacute and post-acute phases of stroke. For over more than two decades, therapeutic efforts in the stroke field have focused on the promotion of neuronal survival, which failed to succeed in clinical trials in humans until now (Savitz and Fisher, 2007; Ginsberg, 2008). From failure to translate successful neuroprotection therapies from animal models to humans, it may be concluded that the stimulation of survival alone is without prospect, as long as no successful remodeling of brain tissue stimulated by a permissive microenvironment takes place. Indeed, studies done recently have shown that extensive remodeling occurs in the brain following an ischemic event (Hermann and Zechariah, 2009; Hermann and Chopp, 2012). Currently, there is hope that stroke recovery might be promoted through pharmacological or cell-based therapies. Indeed, promising results from experimental studies have led to clinical trials, the results of which are currently awaited (Lener et al., 2015).

Remodeling of ischemic brain tissue involves interactions between neurons, glial and microvascular cells that create a microenvironment in which neurological recovery may ensue. Neurons and brain capillaries sprout. Neuronal outgrowth enables the formation of functional axons and synapses in the brain both over long [e.g., along pyramidal tract (Andres et al., 2011; Reitmeir et al., 2011)] and short (e.g., within motor cortex (Clarkson et al., 2010; Hermann and ElAli, 2012)) distances, thus allowing for the restitution of neuronal networks that were damaged by the stroke. The remodeling of ischemic brain tissue also includes responses of immature cells, namely of endothelial progenitor cells (EPC), neural progenitor cells (NPC), and inflammatory cells. New blood vessels are formed, and EPC and NPC are attracted to the stroke lesion. Glial cells contribute to the remodeling of the extracellular matrix, enabling neuronal plasticity.

In the process of brain remodeling, proliferating microvascular cells play a supportive role, enabling the migration of neural precursor cells and promoting the remodeling of neurons and glial cells via secretion of growth factors (Hermann and Zechariah, 2009). This rearrangement of cell-cell interactions is followed by the recovery of the BBB, leading to the restoration of brain homeostasis (Hermann and ElAli, 2012).

The remodeling potential of the NVU serves as an important therapeutic target in stroke and other acute neurologic conditions. After stroke, damaged and inflamed endothelium release pro-inflammatory factors and extracellular vesicles (EVs) that pass through the leaky BBB and activate astrocytes and microglia to release pro-inflammatory cytokines (TNF $\alpha$ , IL-1 $\beta$ ) (Norden et al., 2014). Microglia also release the anti-inflammatory cytokine IL-10 that acts on reactive astrocytes to modify their cytokine secretion from a pro-inflammatory profile toward a pro-recovery one, represented mainly by TGF $\beta$ . During the post-stroke BBB repair and parenchymal remodeling process, NVU cells cooperate and release pro-recovery factors (e.g., IL-4, IL-10, TGF $\beta$ ) that switch microglia into a pro-remodeling phenotype that release growth factors (Norden et al., 2014). Moreover, IL-10 acts on the endothelial and vascular cells to modulate vascular repair and remodeling, diminishes leukocyte–endothelial interactions, decreases expression and activation of cytokine receptors, promotes NO-induced vasodilatation and diminish ROS production and oxidative stress by inhibiting a NADPH oxidase subunit (Nox1) with impact on degenerative vascular remodeling (Dammanahalli et al., 2011; Garcia et al., 2017). Overall, IL-10 secreted from the NVU cells, but also from MSCs and their exosomes (Nakajima et al., 2017), is a pro-survival factor for neurons and glial cells that diminishes the post-lesional inflammatory response and limits the secondary damage during the resolution phase (Mosser and Zhang, 2008).

Pericytes behavior in different phases of ischemic stroke were recently described (Yang et al., 2017). Briefly, during the hyperacute phase, pericytes constriction causes capillary occlusion (no-reflow phenomenon). Then, during the acute phase, pericytes have a pro-inflammatory and immune-modulatory action, with consecutive increase in BBB permeability and brain edema. By protecting the endothelium on its abluminal side and through release of neurotrophins, pericytes stabilize the BBB and protect brain parenchyma. Further, during post-stroke recovery phase, pericytes have a neuroprotective activity, promoting angiogenesis, neurogenesis, and brain recovery. The complex multifaceted, multistage pericytes intervention in ischemic-reperfusion injury and repair processes, recommend them for new targeted therapeutic strategies (Cai et al., 2017).

## Neurovascular Unit-Derived Exosomes in Response to Stroke

Pericytes are important players in post-stroke NVU remodeling. Thus, they were shown to become activated and gain multipotent stem cell phenotype after brain ischemia and express the neuroepithelial stem cell marker nestin, with a potential to differentiate into neural and vascular precursor lineages (Nakagomi et al., 2015). Cooperation between endothelial cells and pericytes occurs both through paracrine interaction, but also through an exosomal bidirectional communication and is essential for preserving the microvascular functionality and stability. For example, endothelium or pericyte-derived hypoxic exosomes were shown to induce an angiogenic program (Fan, 2014; Mayo and Bearden, 2015). Secondary to local injury and perivascular inflammation, MSCs are released from their perivascular location and secrete bioactive molecules and exosomes with immunomodulatory and trophic effects, supporting the regenerative microenvironment needed for the post-injury recovery (Caplan and Correa, 2011). A recent work even suggests that perivascular MSCs are adventitial cells, acting as precursors of pericytes and other stromal cells during tissue homeostasis (de Souza et al., 2016). Not all pericytes can generate MSCs. It has been recently shown that from the various sub-population of existing pericytes, not all of them can act like stem cells, and some act like fibroblasts.

Neuronal exosomes are present at synaptic level, both within pre- and postsynaptic compartments, and transport synaptic receptors (e.g., AMPA receptors, GPCRs) (Koniusz et al., 2016) contributing to synaptic plasticity, both locally and within broader neuronal networks (Chen and Chopp, 2018). The activity within glutamatergic synapses, which is increased in post-stroke excitotoxic conditions, stimulates neuronal release of exosomes that preferentially bind to adjacent neurons, impacting on interneuronal communication (Chivet et al., 2014). The exosomes released secondary to neuronal depolarization are rich in miRNAs, potentially promoting synaptic plasticity by enabling the rapid translation of associated proteins (Goldie et al., 2014).

The interactions between neurons, glial cells and microvascular cells are finely tuned. They involve mutual cell to cell communication via release of growth factors as well as physical cell-cell interactions across the extracellular matrix that is itself subjected to remodeling processes after stroke (Rosell and Lo, 2008). Considering the complexity of these systems and considering both the structural and functional heterogeneities of brain structures and the heterogeneities of ischemic strokes with regard to their size, etiology, and localization (Hermann and Chopp, 2012), the development of neurorestorative therapies is a true challenge (Hermann et al., 2015).

in Response to

Stroke

## The Potential Use of Mesenchymal Stem Cells and Their Exosomes for Stroke Therapy

Mesenchymal Stem

Cells and Their

Exosomes for Stroke

Ischemic stroke is a leading cause of death and long-term disability in industrialized countries, with thrombolysis and

interventional vascular recanalization being the only treatments available. Due to severe side effects and a narrow

therapeutic time window, only a small proportion of stroke patients receive this therapy. Thus, additional therapeutic

concepts are mandatory (Hermann and Chopp, 2012). Strategies that promote neuronal survival in the acute stroke

phase have successfully been studied in experimental stroke models, but were not successful in clinical trials. Therefore,

the research focus has recently shifted from the acute to post-acute stroke phase (Hermann and Chopp, 2012). After

acknowledging that transplanted cells integrate poorly into existing neural networks and that they induce brain

remodeling in a paracrine way by secreting a heterogeneous group of nanovesicles, these EVs have been identified as key

players that mediate restorative effects of stem and progenitor cells in ischemic brain tissue. Neuroprotection as

observed after EV infusion in experimental stroke models is related to stem cell application in stroke. As a matter of fact,

stem cell-induced neurological recovery after stroke is not a consequence of cell regeneration but due to paracrine

mechanisms of grafted cells, among which stem cell-derived EVs are key mediators (Doeppner et al., 2018).

Blood-brain barrier may block or diminish the access of therapeutic agents within the central nervous system and therefore many nervous system diseases lack an efficient treatment because of a deficient drug delivery vehicle.

Considering this important issue, research is nowadays developing nanocarriers for brain targeted drug delivery and exploit the potential use of stem cells to secrete exosomes, as natural nanovesicles rich in biological active molecules. As lipid-bound nanoparticles, exosomes easily interconnect cells and cross selective-permeable membranes such as BBB, thus emerging as versatile tools for new therapeutic strategies (e.g., regenerative, immune-modulatory or anti-tumor therapies), either acting through their biochemically active constituents (e.g., proteins, lipids, genetic material), or serving as natural nonimmunogenic vehicles for drug delivery (Lener et al., 2015).

## Non-exosomal Effects of Mesenchymal Stem Cells

The human brain contains reservoirs of neural stem and precursor cells in the subventricular zone (SVZ) surrounding the lateral ventricles (Bacigaluppi et al., 2009). Although cerebral ischemia triggers the activation of these cells and promotes their migration toward ischemic lesion sites, their siblings hardly survive and differentiate within the ischemic milieu (Doeppner et al., 2012, 2014b). To improve brain remodeling and plasticity, and to bypass limitations of endogenous neurogenesis following ischemic stroke, a variety of approaches started to focus on the transplantation of NPCs or somatic stem cell entities, such as MSCs (Popa-Wagner et al., 2006, 2007, 2011). MSCs secretome comprises growth factors, cytokines, chemokines, extracellular matrix components, genetic material, but also EVs (exosomes and microvesicles), recommending them as versatile tools in clinical applications (Gaceb et al., 2018).

Mesenchymal stem cells, known as “sentinel and safe-guards of injury” (Caplan, 2016), were shown to produce neurotrophic factors such as nerve growth factor, brain-derived neurotrophic factor, or glial-derived neurotrophic factor, explaining their therapeutic potential (Lopatina et al., 2011).

**TABLE OF CONTENTS**  
Initially, it was assumed that transplanted NPCs and MSCs home to affected sites and, upon expansion and differentiation, directly replace the lost brain cells to restore tissue functions. In this context, our lab has comprehensively characterized the therapeutic effects of SVZ-derived adult NPCs in a mouse model of ischemic stroke, i.e., transient proximal (i.e., intraluminal) middle cerebral artery occlusion (MCAO). We observed that systemic NPC delivery induces profound brain tissue remodeling, reflected by reduced secondary neurodegeneration, reduced neuroinflammation, reduced astrogliosis and reduced microglial activation, that was associated with functional neurological recovery (Bacigaluppi et al., 2009; Doeppner et al., 2012, 2014a,b). Remarkably, it turned out that systemic intravenous administration of adult NPCs was more effective than intracerebral transplantation. Indeed, systemic administration effectively resulted in the stabilization of BBB integrity. However, just 0.1–0.3% of intravenously transplanted NPCs were detected in the brain and most of them were in an undifferentiated state (Bacigaluppi et al., 2009). These findings imply that NPCs act in a paracrine rather than a cellular mode.

Derived Exosomes  
in Response to

## Exosomal Effects of Mesenchymal Stem Cells

The Potential Use of Exosomes  
Cells and Their Exosomes for Stroke Therapy  
Conclusion and Perspectives  
Author  
Conflict of Interest  
MSCs act after an ischemic stroke. Thus, MSCs overexpressing IL-10 improved neuronal survival in the ischemic hemisphere (Nakajima et al., 2017). Interestingly, MSCs-derived exosomes were shown to exhibit post-stroke changes in their miRNA profile, mostly in the miRNAs actively involved in the repair process by altering gene expression and promoting brain recovery (Liu et al., 2013).

Classically, paracrine effects were thought to be mediated by soluble molecules such as growth factors, cytokines, chemokines and hormones. More recent data, however, demonstrate that several physiological and pathophysiological processes are controlled by exosomes (Cramer et al., 2017). In experimental stroke models, evidence was provided that exosomes exert neuroprotective, proangiogenic and neuronal plasticity-promoting functions (Xin et al., 2013). Thus, systemic administration of MSC-derived exosomes in a rat model of stroke improved functional recovery and enhanced neurite remodeling, neurogenesis, and angiogenesis (Xin et al., 2013). Furthermore, administration of combined xenogenic (from mini-pig) adipose-derived mesenchymal stem cell (ADMSC) and ADMSC-derived exosome therapy has been shown to reduce brain-infarct zone (BIZ) and enhance neurological recovery in rat after acute ischemic stroke (Chen et al., 2016). At molecular level the beneficial effects of MSC-derived exosomes could be mediated by the miR-17-92 cluster. Thus, rats subjected MCAO and treated with miR-17-92 cluster-enriched exosomes, performed significantly better than the control rats treated with MSC exosome alone (Xin et al., 2017a). Similarly, administration of exosomes isolated from miR-133b-overexpressing MSCs lead to increased neural plasticity and improvement of functional recovery after stroke in rats (Xin et al., 2017b).

Based on these observations, we performed a direct head-by-head comparison of the therapeutic effects of MSCs and their exosomes in a murine model of transient intraluminal MCAO, which predominantly affects the striatum and most lateral parts of the overlying cerebral cortex, showing that systemic MSCs and MSC-derived exosomes are equally effective in enhancing stroke-related motor and coordination recovery thereby confirming the beneficial effects of the exosome therapy reported by Chopp and colleagues who observed a significant reduction in neurological impairment that improved gradually over 4 weeks after systemic delivery of MSC-derived EVs (MSC-EVs) in a model of transient MCAO in rats (Xin et al., 2013). Both therapies promoted post-ischemic endogenous neurogenesis and angiogenesis and reversed the stroke-associated immunodepression (Doeppner et al., 2015).

## Conclusion and Perspectives

Ischemic stroke is a leading cause of death and long-term disability for which no restorative therapy is available. After stroke, the NVU is compromised and has become a major target for restorative therapies in the central nervous system. Emerging research has revealed that the nano-sized exosomes could be used for the NVU remodeling after stroke, due to their ability to mediate cell-to-cell communication. Considering the side effects typically attributed to cell-based therapies, in particular, malignant transformation of the transplanted cells, MSC-derived exosomes are attractive candidates for stroke therapy, as emphasized by a recent position paper (Lener et al., 2015). Indeed, systemic

administration of MSC-derived exosomes is effective in enhancing stroke-related motor and coordination recovery in experimental stroke models, fueling the hope for clinical studies. Nevertheless, for clinical applications we need further studies to shed light on (i) mechanisms of the interaction between exosomes and target cells, (ii) circulation kinetics and biodistribution; (iii) biogenesis mechanism; (iv) potential side effects. For example, tumor-secreted exosomes may act as mediators in cancer metastasis by maintenance and enhancement of tumor microenvironment (Salido-Guadarrama et al., 2014; Cheng et al., 2017; Li et al., 2018). Likewise, several studies have reported high levels of cholesterol ester (CE), triacylglycerol (TAG) and cardiolipin in exosomal preparations fueling concerns about increasing the risk of stroke instead of having a beneficial effect (Llorente et al., 2007; Van Meer et al., 2008; Strauss et al., 2010; Record et al., 2014; Sklindas et al., 2017; Popa-Wagner et al., 2018).

Furthermore, considering that ischemic stroke mainly affects elderly patients, experimental data in aged rodents are urgently required before a clinical proof-of-concept study in human patients should be envisaged (Popa-Wagner et al., 2006, 2007, 2018; Balseanu et al., 2014).

Exosomes for Stroke

Therapy

## Author Contributions

Concussion and

Perspectives

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

Contributions

Funding

## Funding

Interest

Statement

This work was supported by grants of the Romanian National Authority for Science Research and Innovation, CNCS –

Footnotes

UEFISCDI, project numbers PN-III-P4-ID-PCE-2016-0340; PN-III-P2-2.1-PED-2016-1013 and PN-III-P4-ID-PCE-

References

2016-0215.

## Conflict of Interest Statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Footnotes

1. ^<http://www.exocarta.org/> (<http://www.exocarta.org/>)

## References

- Abbott, N. J. (2002). Astrocyte–endothelial interactions and blood–brain barrier permeability. *J. Anat.* 200, 629–638. doi: 10.1046/j.1469-7580.2002.00064.x
- [CrossRef Full Text](https://doi.org/10.1046/j.1469-7580.2002.00064.x) ([Google Scholar](http://scholar.google.com/scholar_lookup?&title=Astrocyte-endothelial+interactions+and+blood-brain+barrier+permeability%2E&journal=J%2E+Anat%2E&author=Abbott+N.+J.&publication_year=2002&volume=200&pages=629-638))
- Abbott, N. J. (2013). Blood–brain barrier structure and function and the challenges for CNS drug delivery. *J. Inherit. Metab. Dis.* 36, 437–449. doi: 10.1007/s10545-013-9608-0
- [PubMed Abstract](https://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=23609350) ([CrossRef Full Text](https://doi.org/10.1007/s10545-013-9608-0)) ([Google Scholar](http://scholar.google.com/scholar_lookup?&title=Blood-brain+barrier+structure+and+function+and+the+challenges+for+CNS+drug+delivery%2E&journal=J%2E+Inherit%2E+Metab%2E+Dis%2E&author=Abbott+N.+J.&publication_year=2013&volume=36&pages=437-449))
- Abbott, N. J., and Friedman, A. (2012). Overview and introduction: the blood-brain barrier in health and disease. *Epilepsia* 53(Suppl. 6), 1–6. doi: 10.1111/j.1528-1167.2012.03696.x
- [PubMed Abstract](https://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=23134489) ([CrossRef Full Text](https://doi.org/10.1007/s10545-013-9608-0)) ([Google Scholar](http://scholar.google.com/scholar_lookup?&title=Overview+and+introduction%3A+the+blood-brain+barrier+in+health+and+disease%2E&journal=Epilepsia&author=Abbott+N.+J.&author=and+Friedman+A.&publication_year=2012&pages=1-6))
- András, I. E., and Toborek, M. (2016). Extracellular vesicles of the blood-brain barrier. *Tissue Barriers* 4:e1131804. doi: 10.1080/21688370.2015.1131804
- [PubMed Abstract](https://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=27141419) ([CrossRef Full Text](https://doi.org/10.1080/21688370.2015.1131804)) ([Google Scholar](http://scholar.google.com/scholar_lookup?&title=Exosomes+as+a+biological+tool+for+extracellular+vesicles+of+the+blood-brain+barrier%2E&journal=Tissue+Barriers&author=András+I.+E.&author=and+Toborek+M.&publication_year=2016))
- Andres, R. H., Horie, N., Slikker, W., Keren-Gill, H., Zhan, K., Sun, G., et al. (2011). Human neural stem cells enhance structural plasticity and axonal transport in the ischaemic brain. *Brain* 134, 1777–1789. doi: 10.1093/brain/awr094
- Remodeling in response to stroke. [PubMed Abstract](https://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=21616972) ([CrossRef Full Text](https://doi.org/10.1093/brain/awr094)) ([Google Scholar](http://scholar.google.com/scholar_lookup?&title=Human+neural+stem+cells+enhance+structural+plasticity+and+axonal+transport+in+the+ischaemic+brain%2E&journal=Brain&author=Andres+R.+H.&author=Horie+N.&author=Slikker+W.&author=Keren-Gill+H.&author=Zhan+K.&author=Sun+G.&publication_year=2011&volume=134&pages=1777-1789))
- Attwell, D., Buchan, A. M., Charpak, S., Lauritzen, M., Macvicar, B. A., and Newman, E. A. (2010). Glial and neuronal control of brain blood flow. *Nature* 468, 232–243. doi: 10.1038/nature09613
- The Potential Use of Mesenchymal Stem Cells in Therapy for Stroke. [PubMed Abstract](https://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=21068832) ([CrossRef Full Text](https://doi.org/10.1038/nature09613)) ([Google Scholar](http://scholar.google.com/scholar_lookup?&title=Glial+and+neuronal+control+of+brain+blood+flow%2E&journal=Nature&author=Attwell+D.&author=Buchan+A.+M.&author=Charpak+S.&author=Lauritzen+M.&author=Macvicar+B.+A.&author=and+Newman+E.+A.&publication_year=2010&volume=468&pages=232-243))
- Bacigaluppi, M., Pluchino, S., Peruzzotti-Jametti, L., Kilic, E., Kilic, U., Salani, G., et al. (2009). Delayed post-ischaemic neuroprotection following systemic neural stem cell transplantation involves multiple mechanisms. *Brain* 132, 2239–2251. doi: 10.1093/brain/awp174
- Perspectives. [PubMed Abstract](https://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=19617198) ([CrossRef Full Text](https://doi.org/10.1093/brain/awp174)) ([Google Scholar](http://scholar.google.com/scholar_lookup?&title=Delayed+post-ischaemic+neuroprotection+following+systemic+neural+stem+cell+transplantation+involves+multiple+mechanisms%2E&journal=Brain&author=Bacigaluppi+M.&author=Pluchino+S.&author=Peruzzotti-Jametti+L.&author=Kilic+E.&author=Kilic+U.&author=Salani+G.&publication_year=2009&volume=132&pages=2239-2251))
- Balseanu, A. T., Buga, A. M., Catalin, B., Wagner, D. C., Boltze, J., Zagrean, A. M., et al. (2014). Multimodal approaches for regenerative stroke therapies: combination of granulocyte colony-stimulating factor with bone marrow mesenchymal stem cells is not superior to G-CSF alone. *Front. Aging Neurosci.* 6:130. doi: 10.3389/fnagi.2014.00130
- Footnotes. [PubMed Abstract](https://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=25002846) ([CrossRef Full Text](https://doi.org/10.3389/fnagi.2014.00130)) ([Google Scholar](http://scholar.google.com/scholar_lookup?&title=Multimodal+approaches+for+regenerative+stroke+therapies%3A+combination+of+granulocyte+colony-stimulating+factor+with+bone+marrow+mesenchymal+stem+cells+is+not+superior+to+G-CSF+alone%2E&journal=Front%2E+Aging+Neurosci%2E&author=Balseanu+A.+T.&author=Buga+A.+M.&author=Catalin+B.&author=Wagner+D.+C.&author=Boltze+J.&author=Zagrean+A.+M.&publication_year=2014))
- Balusu, S., Van Wonterghem, E., De Rycke, R., Raemdonck, K., Stremersch, S., Gevaert, K., et al. (2016). Identification of a novel mechanism of blood–brain communication during peripheral inflammation via choroid plexus derived extracellular vesicles. *EMBO Mol. Med.* 8, 1162–1183. doi: 10.15252/emmm.201606271
- [PubMed Abstract](https://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=27596437) ([CrossRef Full Text](https://doi.org/10.15252/emmm.201606271)) ([Google Scholar](http://scholar.google.com/scholar_lookup?&title=Identification+of+a+novel+mechanism+of+blood-brain+communication+during+peripheral+inflammation+via+choroid+plexus+derived+extracellular+vesicles%2E&journal=EMBO+Mol%2E++Med%2E&author=Balusu+S.&author=Van+Wonterghem+E.&author=De+Rycke+R.&author=Raemdonck+K.&author=Stremersch+S.&author=Gevaert+K.&publication_year=2016&volume=8&pages=1162-1183))
- Bang, C., and Thum, T. (2012). Exosomes: new players in cell-cell communication. *Int. J. Biochem. Cell Biol.* 44, 2060–2064. doi: 10.1016/j.biocel.2012.08.007

[PubMed Abstract](https://pubmed.ncbi.nlm.nih.gov/22903023/) (<https://doi.org/10.1016/j.biocel.2012.08.007>) | [CrossRef Full Text](https://scholar.google.com/scholar_lookup?&title=Exosomes%3A+new+players+in+cell-cell+communication%2E&journal=Int%2E+J%2E+Biochem%2E+Cell+BioI%2E&author=Bang+C.&author=and+Thum+T.&publication_year=2012&volume=44&pages=2060-2064)

Barile, L., and Vassalli, G. (2017). Exosomes: therapy delivery tools and biomarkers of diseases. *Pharmacol. Ther.* 174, 63–78. doi: [10.1016/j.pharmthera.2017.02.020](https://doi.org/10.1016/j.pharmthera.2017.02.020)

[PubMed Abstract](https://pubmed.ncbi.nlm.nih.gov/28202367/) (<https://doi.org/10.1016/j.pharmthera.2017.02.020>) | [CrossRef Full Text](https://scholar.google.com/scholar_lookup?&title=Exosomes%3A+therapy+delivery+tools+and+biomarkers+of+diseases%2E&journal=Pharmacol%2E+Ther%2E&author=Barile+L.&author=and+Vassalli+G.&publication_year=2017&volume=174&pages=63-78)

Bellingham, S. A., Guo, B. B., Coleman, B. M., and Hill, A. F. (2012). Exosomes: vehicles for the transfer of toxic proteins associated with neurodegenerative diseases? *Front. Physiol.* 3:124. doi: [10.3389/fphys.2012.00124](https://doi.org/10.3389/fphys.2012.00124)

[PubMed Abstract](https://pubmed.ncbi.nlm.nih.gov/22563321/) (<https://doi.org/10.3389/fphys.2012.00124>) | [CrossRef Full Text](https://scholar.google.com/scholar_lookup?&title=Exosomes%3A+vehicles+for+the+transfer+of+toxic+proteins+associated+with+neurodegenerative+diseases%2E&journal=Front%2E+Physiol%2E&author=Bellingham+S.+A.&author=Guo+B.+B.&author=Coleman+B.+M.&author=and+Hill+A.+F.&publication_year=2012)

Burrello, J., Monticone, S., Gai, C., Gomez, Y., Kholia, S., and Camussi, G. (2016). Stem cell-derived extracellular vesicles and immune-modulation. *Front. Cell Dev. Biol.* 4:83. doi: [10.3389/fcell.2016.00083](https://doi.org/10.3389/fcell.2016.00083)

**TABLE OF CONTENTS**  
[CrossRef Full Text](https://doi.org/10.3389/fcell.2016.00083) (<https://doi.org/10.3389/fcell.2016.00083>) | [Google Scholar](https://scholar.google.com/scholar_lookup?&title=Stem+cell-derived+extracellular+vesicles+and+immune-modulation%2E&journal=Front%2E+Cell+Dev%2E+BioI%2E&author=Burrello+J.&author=Monticone+S.&author=Gai+C.&author=Gomez+Y.&author=Kholia+S.&author=and+Camussi+G.&publication_year=2016) ([https://scholar.google.com/scholar\\_lookup?&title=Stem+cell-derived+extracellular+vesicles+and+immune-modulation%2E&journal=Front%2E+Cell+Dev%2E+BioI%2E&author=Burrello+J.&author=Monticone+S.&author=Gai+C.&author=Gomez+Y.&author=Kholia+S.&author=and+Camussi+G.&publication\\_year=2016](https://scholar.google.com/scholar_lookup?&title=Stem+cell-derived+extracellular+vesicles+and+immune-modulation%2E&journal=Front%2E+Cell+Dev%2E+BioI%2E&author=Burrello+J.&author=Monticone+S.&author=Gai+C.&author=Gomez+Y.&author=Kholia+S.&author=and+Camussi+G.&publication_year=2016))

Cai, W., Liu, H., Zhao, J., Chen, L. Y., Chen, J., Lu, Z., et al. (2017). Pericytes in brain injury and repair after ischemic stroke. *Trans. Stroke Res.* 8, 107–121. doi: [10.1007/s12975-016-0504-4](https://doi.org/10.1007/s12975-016-0504-4)

Biological

[PubMed Abstract](https://pubmed.ncbi.nlm.nih.gov/27837475/) (<https://doi.org/10.1007/s12975-016-0504-4>) | [Google Scholar](https://scholar.google.com/scholar_lookup?&title=Pericytes+in+brain+injury+and+repair+after+ischemic+stroke%2E&journal=Trans%2E+Stroke+Res%2E&author=Cai+W.&author=Liu+H.&author=Zhao+J.&author=Chen+L.+Y.&author=Chen+J.&author=Lu+Z.&publication_year=2017&volume=8&pages=107-121) ([https://scholar.google.com/scholar\\_lookup?&title=Pericytes+in+brain+injury+and+repair+after+ischemic+stroke%2E&journal=Trans%2E+Stroke+Res%2E&author=Cai+W.&author=Liu+H.&author=Zhao+J.&author=Chen+L.+Y.&author=Chen+J.&author=Lu+Z.&publication\\_year=2017&volume=8&pages=107-121](https://scholar.google.com/scholar_lookup?&title=Pericytes+in+brain+injury+and+repair+after+ischemic+stroke%2E&journal=Trans%2E+Stroke+Res%2E&author=Cai+W.&author=Liu+H.&author=Zhao+J.&author=Chen+L.+Y.&author=Chen+J.&author=Lu+Z.&publication_year=2017&volume=8&pages=107-121))

Biomedeling in

Caplan, A. I. (2008). All MSCs are pericytes? *Cell Stem Cell* 3, 229–230. doi: [10.1016/j.stem.2008.08.008](https://doi.org/10.1016/j.stem.2008.08.008)

Response to Stroke

[PubMed Abstract](https://pubmed.ncbi.nlm.nih.gov/18786406/) (<https://doi.org/10.1016/j.stem.2008.08.008>) | [Google Scholar](https://scholar.google.com/scholar_lookup?&title=All+MSCs+are+pericytes%2E&journal=Cell+Stem+Cell&author=Caplan+A.+I.&publication_year=2008&volume=3&pages=229-230) ([https://scholar.google.com/scholar\\_lookup?&title=All+MSCs+are+pericytes%2E&journal=Cell+Stem+Cell&author=Caplan+A.+I.&publication\\_year=2008&volume=3&pages=229-230](https://scholar.google.com/scholar_lookup?&title=All+MSCs+are+pericytes%2E&journal=Cell+Stem+Cell&author=Caplan+A.+I.&publication_year=2008&volume=3&pages=229-230))

in Response to

Caplan, A. I. (2016). MSCs: the sentinel and safe-guards of injury. *J. Cell. Physiol.* 231, 1413–1416. doi: [10.1002/jcp.25255](https://doi.org/10.1002/jcp.25255)

Stroke

[PubMed Abstract](https://pubmed.ncbi.nlm.nih.gov/26565391/) (<https://doi.org/10.1002/jcp.25255>) | [Google Scholar](https://scholar.google.com/scholar_lookup?&title=MSCs%3A+the+sentinel+and+safe-guards+of+injury%2E&journal=J%2E+Cell%2E++Physiol%2E&author=Caplan+A.+I.&publication_year=2016&volume=231&pages=1413-1416) ([https://scholar.google.com/scholar\\_lookup?&title=MSCs%3A+the+sentinel+and+safe-guards+of+injury%2E&journal=J%2E+Cell%2E++Physiol%2E&author=Caplan+A.+I.&publication\\_year=2016&volume=231&pages=1413-1416](https://scholar.google.com/scholar_lookup?&title=MSCs%3A+the+sentinel+and+safe-guards+of+injury%2E&journal=J%2E+Cell%2E++Physiol%2E&author=Caplan+A.+I.&publication_year=2016&volume=231&pages=1413-1416))

Cells and Their

Caplan, A. I., and Correa, D. (2011). The MSC: an injury drugstore. *Cell Stem Cell* 9, 11–15. doi: [10.1016/j.stem.2011.06.008](https://doi.org/10.1016/j.stem.2011.06.008)

Exosomes for Stroke

[PubMed Abstract](https://pubmed.ncbi.nlm.nih.gov/21726829/) (<https://doi.org/10.1016/j.stem.2011.06.008>) | [Google Scholar](https://scholar.google.com/scholar_lookup?&title=The+MSC%3A+an+injury+drugstore%2E&journal=Cell+Stem+Cell&author=Caplan+A.+I.&author=and+Correa+D.&publication_year=2011&volume=9&pages=11-15) ([https://scholar.google.com/scholar\\_lookup?&title=The+MSC%3A+an+injury+drugstore%2E&journal=Cell+Stem+Cell&author=Caplan+A.+I.&author=and+Correa+D.&publication\\_year=2011&volume=9&pages=11-15](https://scholar.google.com/scholar_lookup?&title=The+MSC%3A+an+injury+drugstore%2E&journal=Cell+Stem+Cell&author=Caplan+A.+I.&author=and+Correa+D.&publication_year=2011&volume=9&pages=11-15))

Perspectives

Chen, J., and Chopp, M. (2018). Exosome therapy for stroke. *Stroke* 49, 1083–1090. doi: [10.1161/STROKEAHA.117.018292](https://doi.org/10.1161/STROKEAHA.117.018292)

Author

[PubMed Abstract](https://pubmed.ncbi.nlm.nih.gov/29669873/) (<https://doi.org/10.1161/STROKEAHA.117.018292>) | [Google Scholar](https://scholar.google.com/scholar_lookup?&title=Exosome+therapy+for+stroke%2E&journal=Stroke&author=Chen+J.&author=and+Chopp+M.&publication_year=2018&volume=49&pages=1083-1090) ([https://scholar.google.com/scholar\\_lookup?&title=Exosome+therapy+for+stroke%2E&journal=Stroke&author=Chen+J.&author=and+Chopp+M.&publication\\_year=2018&volume=49&pages=1083-1090](https://scholar.google.com/scholar_lookup?&title=Exosome+therapy+for+stroke%2E&journal=Stroke&author=Chen+J.&author=and+Chopp+M.&publication_year=2018&volume=49&pages=1083-1090))

Conflict of Interest

Chen, K.-H., Chen, C.-H., Wallace, C. G., Yuen, C.-M., Kao, G.-S., Chen, Y.-L., et al. (2016). Intravenous administration of xenogenic adipose-derived mesenchymal stem cells (ADMSC) and ADMSC-derived exosomes markedly reduced brain infarct volume and preserved neurological function in rat after focal ischemic stroke. *Oncotarget* 7, 74537–74556. doi: [10.18632/oncotarget.12902](https://doi.org/10.18632/oncotarget.12902)

[PubMed Abstract](https://pubmed.ncbi.nlm.nih.gov/27793019/) (<https://doi.org/10.18632/oncotarget.12902>) | [Google Scholar](https://scholar.google.com/scholar_lookup?&title=Intravenous+administration+of+xenogenic+adipose-derived+mesenchymal+stem+cells+%28ADMSC%29+and+ADMSC-derived+exosomes+markedly+reduced+brain+infarct+volume+and+preserved+neurological+function+in+rat+after+acute+ischemic+stroke%2E&journal=Oncotarget&author=Chen+K.-H.&author=Chen+C.-H.&author=Wallace+C.+G.&author=Yuen+C.-M.&author=Kao+G.-S.&author=Chen+Y.-L.&publication_year=2016&volume=7&pages=74537-74556) ([https://scholar.google.com/scholar\\_lookup?&title=Intravenous+administration+of+xenogenic+adipose-derived+mesenchymal+stem+cells+%28ADMSC%29+and+ADMSC-derived+exosomes+markedly+reduced+brain+infarct+volume+and+preserved+neurological+function+in+rat+after+acute+ischemic+stroke%2E&journal=Oncotarget&author=Chen+K.-H.&author=Chen+C.-H.&author=Wallace+C.+G.&author=Yuen+C.-M.&author=Kao+G.-S.&author=Chen+Y.-L.&publication\\_year=2016&volume=7&pages=74537-74556](https://scholar.google.com/scholar_lookup?&title=Intravenous+administration+of+xenogenic+adipose-derived+mesenchymal+stem+cells+%28ADMSC%29+and+ADMSC-derived+exosomes+markedly+reduced+brain+infarct+volume+and+preserved+neurological+function+in+rat+after+acute+ischemic+stroke%2E&journal=Oncotarget&author=Chen+K.-H.&author=Chen+C.-H.&author=Wallace+C.+G.&author=Yuen+C.-M.&author=Kao+G.-S.&author=Chen+Y.-L.&publication_year=2016&volume=7&pages=74537-74556))

Cheng, L., Zhang, K., Wu, S., Cui, M., and Xu, T. (2017). Focus on mesenchymal stem cell-derived exosomes: opportunities and challenges in cell-free therapy. *Stem Cells Int.* 2017:6305295. doi: [10.1155/2017/6305295](https://doi.org/10.1155/2017/6305295)

[PubMed Abstract](https://pubmed.ncbi.nlm.nih.gov/29410682/) (<https://doi.org/10.1155/2017/6305295>) | [Google Scholar](https://scholar.google.com/scholar_lookup?&title=Focus+on+mesenchymal+stem+cell-derived+exosomes%3A+opportunities+and+challenges+in+cell-free+therapy%2E&journal=Stem+Cells+Int%2E&author=Cheng+L.&author=Zhang+K.&author=Wu+S.&author=Cui+M.&author=and+Xu+T.&publication_year=2017) ([https://scholar.google.com/scholar\\_lookup?&title=Focus+on+mesenchymal+stem+cell-derived+exosomes%3A+opportunities+and+challenges+in+cell-free+therapy%2E&journal=Stem+Cells+Int%2E&author=Cheng+L.&author=Zhang+K.&author=Wu+S.&author=Cui+M.&author=and+Xu+T.&publication\\_year=2017](https://scholar.google.com/scholar_lookup?&title=Focus+on+mesenchymal+stem+cell-derived+exosomes%3A+opportunities+and+challenges+in+cell-free+therapy%2E&journal=Stem+Cells+Int%2E&author=Cheng+L.&author=Zhang+K.&author=Wu+S.&author=Cui+M.&author=and+Xu+T.&publication_year=2017))

Chivet, M., Javalet, C., Laulagnier, K., Blot, B., Hemming, F. J., and Sadoul, R. (2014). Exosomes secreted by cortical neurons upon glutamatergic synapse activation specifically interact with neurons. *J. Extracell. Vesicles* 3:24722. doi: [10.3402/jev.v3.24722](https://doi.org/10.3402/jev.v3.24722)

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=25398455>) | [CrossRef Full Text](#) (<https://doi.org/10.3402/jev.v3.24722>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup](http://scholar.google.com/scholar_lookup))  
**&title=**Exosomes+secreted+by+cortical+neurons+upon+glutamatergic+synapse+activation+specifically+interact+with+neurons%2E&journal=J%2E+Extracell%2E+Vesicles&author=Chivet+M.&author=Javalet+C.&author=Laulagnier+K.&author=Blot+B.&author=Hemming+F.+J.&author=and+Sadoul+R.&publication\_year=2014)

Clarkson, A. N., Huang, B. S., Macisaac, S. E., Mody, I., and Carmichael, S. T. (2010). Reducing excessive GABA-mediated tonic inhibition promotes functional recovery after stroke. *Nature* 468, 305–309. doi: 10.1038/nature09511

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=21048709>) | [CrossRef Full Text](#) (<https://doi.org/10.1038/nature09511>) | [Google Scholar](#) ([&title=Reducing+excessive+GABA-mediated+tonic+inhibition+promotes+functional+recovery+after+stroke%2E&journal=Nature&author=Clarkson+A.+N.&author=Huang+B.+S.&author=Macisaac+S.+E.&author=Mody+I.&author=and+Carmichael+S.+T.&publication\\_year=2010&volume=468&pages=305-309](http://scholar.google.com/scholar_lookup))

Cramer, S. C., Wolf, S. L., Adams, H. P., Chen, D., Dromerick, A. W., Dunning, K. et al. (2017). Stroke recovery and rehabilitation research: issues, opportunities, and the National Institutes of Health StrokeNet. *Stroke* 48, 813–819. doi: 10.1161/STROKEAHA.116.015501

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=28174324>) | [CrossRef Full Text](#) (<https://doi.org/10.1161/STROKEAHA.116.015501>) | [Google Scholar](#) ([&title=Stroke+recovery+and+rehabilitation+research%3A+issues%2C+opportunities%2C+and+the+National+Institutes+of+Health+StrokeNet%2E&journal=Stroke&author=Cramer+S.+C.&author=+Wolf+S.+L.&author=+Adams+H.+P.&author=+Chen+D.&author=+Dromerick+A.+W.&author=+Dunning+K.&author=+Sun+Z.&publication\\_year=2017&volume=48&pages=813-819](http://scholar.google.com/scholar_lookup))

**TABLE OF CONTENT** Wang, X., and Sun, Z. (2011). Genetic interleukin-10 deficiency causes vascular remodeling via the upregulation of Nox1. *J. Hypertens.* 29, 2116–2125. doi: 10.1097/JHH.0b013e32834b22ao

**Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=21918473>) | [CrossRef Full Text](#) (<https://doi.org/10.1097/JHH.0b013e32834b22ao>) | [Google Scholar](#) ([&title=Genetic+interleukin-10+deficiency+causes+vascular+remodeling+via+the+upregulation+of+Nox1%2E&journal=J%2E+Hypertens%2E&author=Dammanahalli+J.+K.&author=+Wang+X.&author=+Sun+Z.&publication\\_year=2011&volume=29&pages=2116-2125](http://scholar.google.com/scholar_lookup))

**Biological Activity** de Souza, L. E., Malta, T. M., Kashima Haddad, S., and Covas, D. T. (2016). Mesenchymal stem cells and pericytes: to what extent are they related? *Stem Cells Dev.* 25, 1843–1852. doi: 10.1089/scd.2016.0109

**Tool** Doeppner, T. R., Bähr, M., Giebel, B., and Hermann, D. M. (2018). Immunological and non-immunological effects of stem cell-derived extracellular vesicles on the ischaemic brain. *Ther. Adv. Neurol. Disord.* 26:1756286418789326. doi: 10.1177/1756286418789326  
**Derived Exosomes**

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=30083231>) | [CrossRef Full Text](#) (<https://doi.org/10.1177/1756286418789326>) | [Google Scholar](#) ([&title=Immunological+and+non-immunological+effects+of+stem+cell-derived+extracellular+vesicles+on+the+ischaemic+brain%2E&journal=Ther%2E++Adv%2E++Neurol%2E++Disord%2E&author=Doeppner+T.+R.&author=Bähr+M.&author=Giebel+B.&author=and+Hermann+D.+M.&publication\\_year=2018](http://scholar.google.com/scholar_lookup))

**Cells and Their Therapeutic Potential** Doeppner, T. R., Ewert, T. A., Tönges, L., Herz, J., Zechariah, A., ElAli, A., et al. (2012). Transduction of neural precursor cells with TAT-Hsp70 chaperone: therapeutic potential against ischemic stroke after intrastratal and systemic transplantation. *Stem Cells* 30, 1297–1310. doi: 10.1002/stem.1098

**Therapy** Doeppner, T. R., Ewert, T. A., Tönges, L., Herz, J., Zechariah, A., ElAli, A., et al. (2012). Transduction of neural precursor cells with TAT-Hsp70 chaperone: therapeutic potential against ischemic stroke after intrastratal and systemic transplantation. *Stem Cells* 30, 1297–1310. doi: 10.1002/stem.1098  
**Perspectives** Doeppner, T. R., Ewert, T. A., Tönges, L., Herz, J., Zechariah, A., ElAli, A., et al. (2012). Transduction of neural precursor cells with TAT-Hsp70 chaperone%3A+therapeutic+potential+against+ischemic+stroke+after+intrastratal+and+systemic+transplantation%2E&journal=Stem+Cells&author=Doeppner+T.+R.&author=Ewert+T.+A.&author=Tönges+L.&author=Herz+J.&author=Zechariah+A.&author=ElAli+A.&publication\_year=2012&volume=30&pages=1297-1310

**Funding** Doeppner, T. R., Herz, J., Görgens, A., Schlechter, J., Ludwig, A.-K., Radtke, S., et al. (2015). Extracellular vesicles improve post-stroke neuroregeneration and prevent postischemic immunosuppression. *Stem Cells Transl. Med.* 4, 1131–1143. doi: 10.5966/sctm.2015-0078

**Conflict of Interest** Doeppner, T. R., Herz, J., Görgens, A., Schlechter, J., Ludwig, A.-K., Radtke, S., et al. (2015). Extracellular vesicles improve post-stroke neuroregeneration and prevent postischemic immunosuppression%2E&journal=Stem+Cells+Transl%2E+Med%2E&author=Doeppner+T.+R.&author=Herz+J.&author=Görgens+A.&author=Schlechter+J.&author=Ludwig+A.-K.  
**Keywords** Radtke+S.&publication\_year=2015&volume=4&pages=1131-1143

Doeppner, T. R., Kaltwasser, B., Bähr, M., and Hermann, D. M. (2014a). Effects of neural progenitor cells on stroke impairment-detailed analysis of behavioral tests. *Front. Cell. Neurosci.* 8:338. doi: 10.3389/fncel.2014.00338

**CrossRef Full Text** (<https://doi.org/10.3389/fncel.2014.00338>) | [Google Scholar](#) ([&title=Effects+of+neural+progenitor+cells+on+stroke+impairment-detailed+analysis+of+behavioral+tests%2E&journal=Front%2E+Cell%2E+Neurosci%2E&author=Doeppner+T.+R.&author=Kaltwasser+B.&author=Bähr+M.&author=and+Hermann+D.+M.&publication\\_year=2014a](http://scholar.google.com/scholar_lookup))

Doeppner, T. R., Kaltwasser, B., Teli, M. K., Bretschneider, E., Bähr, M., and Hermann, D. M. (2014b). Effects of acute versus post-acute systemic delivery of neural progenitor cells on neurological recovery and brain remodeling after focal cerebral ischemia in mice. *Cell Death Dis.* 5:e1386. doi: 10.1038/cddis.2014.359

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=25144721>) | [CrossRef Full Text](#) (<https://doi.org/10.1038/cddis.2014.359>) | [Google Scholar](#) ([&title=Effects+of+acute+versus+post-acute+systemic+delivery+of+neural+progenitor+cells+on+neurological+recovery+and+brain+remodeling+after+focal+cerebral+ischemia+in+mice%2E&journal=Cell+Death+Dis%2E&author=Doeppner+T.+R.&author=Kaltwasser+B.&author=Teli+M.+K.&author=Bretschneider+E.&author=Bähr+M.&author=and+Hermann+D.+M.&publication\\_year=2014b](http://scholar.google.com/scholar_lookup))

We use cookies on this website to improve your user experience. [Learn more \(/www.frontiersin.org/legal/list-of-cookies\)](https://www.frontiersin.org/legal/list-of-cookies)

CLOSE

Dozio, V., and Sanchez, J. C. (2017). Characterisation of extracellular vesicle-subsets derived from brain endothelial cells and analysis of their protein cargo modulation after TNF exposure. *J. Extracell. Vesicles* 6:1302705. doi: 10.1080/20013078.2017.1302705

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=28473883>) | [CrossRef Full Text](#) (<https://doi.org/10.1080/20013078.2017.1302705>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Characterisation+of+extracellular+vesicle-subsets+derived+from+brain+endothelial+cells+and+analysis+of+their+protein+cargo+modulation+after+TNF+exposure%2E&journal=J%2E+Extracel%2E+Vesicles&author=Dozio+V.&author=and+Sanchez+J.+C.&publication\\_year=2017](http://scholar.google.com/scholar_lookup?&title=Characterisation+of+extracellular+vesicle-subsets+derived+from+brain+endothelial+cells+and+analysis+of+their+protein+cargo+modulation+after+TNF+exposure%2E&journal=J%2E+Extracel%2E+Vesicles&author=Dozio+V.&author=and+Sanchez+J.+C.&publication_year=2017))

Fan, G. C. (2014). Hypoxic exosomes promote angiogenesis. *Blood* 124, 3669–3670. doi: 10.1182/blood-2014-10-607846

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=25498451>) | [CrossRef Full Text](#) (<https://doi.org/10.1182/blood-2014-10-607846>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Hypoxic+exosomes+promote+angiogenesis%2E&journal=Blood&author=Fan+G.+C.&publication\\_year=2014&volume=124&pages=3669-3670](http://scholar.google.com/scholar_lookup?&title=Hypoxic+exosomes+promote+angiogenesis%2E&journal=Blood&author=Fan+G.+C.&publication_year=2014&volume=124&pages=3669-3670))

Faure, J., Lachenal, G., Court, M., Hirrlinger, J., Chatellard-Causse, C., Blot, B., et al. (2006). Exosomes are released by cultured cortical neurones. *Mol. Cell. Neurosci.* 31, 642–648. doi: 10.1016/j.mcn.2005.12.003

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=16446100>) | [CrossRef Full Text](#) (<https://doi.org/10.1016/j.mcn.2005.12.003>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Exosomes+are+released+by+cultured+cortical+neurones%2E&journal=Mol%2E+Cell%2E+Neurosci%2E&author=Faure+J.&author=Lachenal+G.&author=Court+M.&author=Hirrlinger+J.&author=Chatellard-Causse+C.&author=Blot+B.&publication\\_year=2006&volume=31&pages=642-648](http://scholar.google.com/scholar_lookup?&title=Exosomes+are+released+by+cultured+cortical+neurones%2E&journal=Mol%2E+Cell%2E+Neurosci%2E&author=Faure+J.&author=Lachenal+G.&author=Court+M.&author=Hirrlinger+J.&author=Chatellard-Causse+C.&author=Blot+B.&publication_year=2006&volume=31&pages=642-648))

Fröhbeis, C., Fröhlich, D., Kuo, W. P., and Krämer-Albers, E. M. (2013). Extracellular vesicles as mediators of neuron-glia communication. *Front. Cell. Neurosci.* 7:182. doi: 10.3389/fncel.2013.00182

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=24194697>) | [CrossRef Full Text](#) (<https://doi.org/10.3389/fncel.2013.00182>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Extracellular+vesicles+as+mediators+of+neuron-glia+communication%2E&journal=Front%2E+Cell%2E++Neurosci%2E&author=Fröhbeis+C.&author=Fröhlich+D.&author=Kuo+W.+P.&author=and+Krämer-Albers+E.+M.&publication\\_year=2013](http://scholar.google.com/scholar_lookup?&title=Extracellular+vesicles+as+mediators+of+neuron-glia+communication%2E&journal=Front%2E+Cell%2E++Neurosci%2E&author=Fröhbeis+C.&author=Fröhlich+D.&author=Kuo+W.+P.&author=and+Krämer-Albers+E.+M.&publication_year=2013))

Biological Communication

Gaceb, A., Barbariga, M., Özén, I., and Paul, G. (2018). The pericyte secretome: potential impact on regeneration. *Biochimie* doi: 10.1016/j.biochi.2018.04.015 [Epub ahead of print].

Tool

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=29698670>) | [CrossRef Full Text](#) (<https://doi.org/10.1016/j.biochi.2018.04.015> [Epub ahead of print].) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=The+pericyte+secretome%3A+potential+impact+on+regeneration%2E&journal=Biochimie&author=Gaceb+A.&author=Barbariga+M.&author=Özén+I.&author=and+Paul+G.&publication\\_year=2018](http://scholar.google.com/scholar_lookup?&title=The+pericyte+secretome%3A+potential+impact+on+regeneration%2E&journal=Biochimie&author=Gaceb+A.&author=Barbariga+M.&author=Özén+I.&author=and+Paul+G.&publication_year=2018))

Garcia, J. M., Stillings, S. A., Leclerc, J. L., Phillips, H., Edwards, N. J., Robicsek, S. A., et al. (2017). Role of interleukin-10 in acute brain injuries. *Front. Neurol.* 8:244. doi: 10.3389/fneur.2017.00244

in Response to Text (<https://doi.org/10.3389/fneur.2017.00244>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Role+of+interleukin-10+in+acute+brain+injuries%2E&journal=Front%2E+Neurol%2E&author=Garcia+J.+M.&author=Stillings+S.+A.&author=Leclerc+J.+L.&author=Phillips+H.&author=Edwards+N.+J.&author=Robicsek+S.+A.&publication\\_year=2017](http://scholar.google.com/scholar_lookup?&title=Role+of+interleukin-10+in+acute+brain+injuries%2E&journal=Front%2E+Neurol%2E&author=Garcia+J.+M.&author=Stillings+S.+A.&author=Leclerc+J.+L.&author=Phillips+H.&author=Edwards+N.+J.&author=Robicsek+S.+A.&publication_year=2017))

The Potential Use of Mesenchymal Stem Cells and Their Exosomes for Stroke

Ginsberg, M. D. (2008). Neuroprotection for ischemic stroke: past, present and future. *Neuropharmacology* 55, 363–389. doi: 10.1016/j.neuropharm.2007.12.007

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=18308347>) | [CrossRef Full Text](#) (<https://doi.org/10.1016/j.neuropharm.2007.12.007>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Neuroprotection+for+ischemic+stroke%3A+past%2C+present+and+future%2E&journal=Neuropharmacology&author=Ginsberg+M.+D.&publication\\_year=2008&volume=55&pages=363-389](http://scholar.google.com/scholar_lookup?&title=Neuroprotection+for+ischemic+stroke%3A+past%2C+present+and+future%2E&journal=Neuropharmacology&author=Ginsberg+M.+D.&publication_year=2008&volume=55&pages=363-389))

Goldie, B. J., Dun, M. D., Lin, M., Smith, N. D., Verrills, N. M., Dayas, C. V., et al. (2014). Activity-associated miRNA are packaged in Map1b-enriched exosomes released from depolarized neurons. *Nucleic Acids Res.* 42, 9195–9208. doi: 10.1093/nar/gku594

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=25053844>) | [CrossRef Full Text](#) (<https://doi.org/10.1093/nar/gku594>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Activity-associated+miRNAs+are+packaged+in+Map1b-enriched+exosomes+released+from+depolarized+neurons%2E&journal=Nucleic+Acids+Res%2E&author=Goldie+B.+J.&author=Dun+M.+D.&author=Lin+N.+D.&author=Smith+N.+D.&author=Verrills+N.+M.&author=Dayas+C.+V.&publication\\_year=2014&volume=42&pages=9195-9208](http://scholar.google.com/scholar_lookup?&title=Activity-associated+miRNAs+are+packaged+in+Map1b-enriched+exosomes+released+from+depolarized+neurons%2E&journal=Nucleic+Acids+Res%2E&author=Goldie+B.+J.&author=Dun+M.+D.&author=Lin+N.+D.&author=Smith+N.+D.&author=Verrills+N.+M.&author=Dayas+C.+V.&publication_year=2014&volume=42&pages=9195-9208))

Footnotes

Goodenough, D. A., Goliger, J. A., and Paul, D. L. (1996). Connexins, connexons, and intercellular communication. *Annu. Rev. Biochem.* 65, 475–502. doi: 10.1146/annurev.bi.65.070196.0002355

**CrossRef Full Text** (<https://doi.org/10.1146/annurev.bi.65.070196.0002355>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Connexins%2C+connexons%2C+and+intercellular+communication%2E&journal=Annu%2E+Rev%2E++Biochem%2E&author=Goodenough+D.+A.&author=Goliger+J.+A.&author=and+Paul+D.+L.&publication\\_year=1996&volume=65&pages=475-502](http://scholar.google.com/scholar_lookup?&title=Connexins%2C+connexons%2C+and+intercellular+communication%2E&journal=Annu%2E+Rev%2E++Biochem%2E&author=Goodenough+D.+A.&author=Goliger+J.+A.&author=and+Paul+D.+L.&publication_year=1996&volume=65&pages=475-502))

Hermann, D., Buga, A. M., and Popa-Wagner, A. (2015). Neurovascular remodeling in the aged ischemic brain. *J. Neural Transm.* 122, S25–S33. doi: 10.1007/s00702-013-1148-0

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=24378703>) | [CrossRef Full Text](#) (<https://doi.org/10.1007/s00702-013-1148-0>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Neurovascular+remodeling+in+the+aged+ischemic+brain%2E&journal=J%2E+Neural+Transm%2E&author=Hermann+D.&author=Buga+A.+M.&author=and+Popa-Wagner+A.&publication\\_year=2015&volume=122&pages=S25-S33](http://scholar.google.com/scholar_lookup?&title=Neurovascular+remodeling+in+the+aged+ischemic+brain%2E&journal=J%2E+Neural+Transm%2E&author=Hermann+D.&author=Buga+A.+M.&author=and+Popa-Wagner+A.&publication_year=2015&volume=122&pages=S25-S33))

Hermann, D. M., and Chopp, M. (2012). Promoting brain remodelling and plasticity for stroke recovery: therapeutic promise and potential pitfalls of clinical translation. *Lancet Neurol.* 11, 369–380. doi: 10.1016/S1474-4422(12)70039-X

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=22441198>) | [CrossRef Full Text](#) ([https://doi.org/10.1016/S1474-4422\(12\)70039-X](https://doi.org/10.1016/S1474-4422(12)70039-X)) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Promoting+brain+remodelling+and+plasticity+for+stroke+recovery%3A+therapeutic+promise+and+potential+pitfalls+of+clinical+translation%2E&journal=Lancet+Neurol%2E&author=Hermann+D.+M.&author=and+Chopp+M.&publication\\_year=2012&volume=11&pages=369-380](http://scholar.google.com/scholar_lookup?&title=Promoting+brain+remodelling+and+plasticity+for+stroke+recovery%3A+therapeutic+promise+and+potential+pitfalls+of+clinical+translation%2E&journal=Lancet+Neurol%2E&author=Hermann+D.+M.&author=and+Chopp+M.&publication_year=2012&volume=11&pages=369-380))

We use cookies on this website to improve your user experience. [Learn more](#) (<https://www.frontiersin.org/legal/list-of-cookies>)

CLOSE

Hermann, D. M., and ElAli, A. (2012). The abluminal endothelial membrane in neurovascular remodeling in health and disease. *Sci. Signal.* 5:re4. doi: 10.1126/scisignal.2002886

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=22871611>) | [CrossRef Full Text](#) (<https://doi.org/10.1126/scisignal.2002886>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=The+abluminal+endothelial+membrane+in+neurovascular+remodeling+in+health+and+disease%2E&journal=Sci%2E++Signal%2E&author=Hermann+D.+M.&author=and+ElAli+A.&publication\\_year=2012](http://scholar.google.com/scholar_lookup?&title=The+abluminal+endothelial+membrane+in+neurovascular+remodeling+in+health+and+disease%2E&journal=Sci%2E++Signal%2E&author=Hermann+D.+M.&author=and+ElAli+A.&publication_year=2012))

Hermann, D. M., and Zechariah, A. (2009). Implications of vascular endothelial growth factor for postischemic neurovascular remodeling. *J. Cereb. Blood Flow Metab.* 29, 1620–1643. doi: 10.1038/jcbfm.2009.100

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=19654590>) | [CrossRef Full Text](#) (<https://doi.org/10.1038/jcbfm.2009.100>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Implications+of+vascular+endothelial+growth+factor+for+postischemic+neurovascular+remodeling%2E&journal=J%2E+Cereb%2E++Blood+Flow+Metabolism%2E&author=Hermann+D.+M.&author=and+Zechariah+A.&publication\\_year=2009&volume=29&pages=1620-1643](http://scholar.google.com/scholar_lookup?&title=Implications+of+vascular+endothelial+growth+factor+for+postischemic+neurovascular+remodeling%2E&journal=J%2E+Cereb%2E++Blood+Flow+Metabolism%2E&author=Hermann+D.+M.&author=and+Zechariah+A.&publication_year=2009&volume=29&pages=1620-1643))

Holm, M. M., Kaiser, J., and Schwab, M. E. (2018). Extracellular vesicles: multimodal envoys in neural maintenance and repair. *Trends Neurosci.* 41, 360–372. doi: 10.1016/j.tins.2018.03.006

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=29605090>) | [CrossRef Full Text](#) (<https://doi.org/10.1016/j.tins.2018.03.006>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Extracellular+vesicles%3A+multimodal+envoys+in+neural+maintenance+and+repair%2E&journal=Trends+Neurosci%2E&author=Holm+M.+M.&author=Kaiser+J.&author=and+Schwab+M.+E.&publication\\_year=2018&volume=41&pages=360-372](http://scholar.google.com/scholar_lookup?&title=Extracellular+vesicles%3A+multimodal+envoys+in+neural+maintenance+and+repair%2E&journal=Trends+Neurosci%2E&author=Holm+M.+M.&author=Kaiser+J.&author=and+Schwab+M.+E.&publication_year=2018&volume=41&pages=360-372))

**TABLE OF CONTENTS**  
Koniusz, S., Andrzejewska, A., Muraca, M., Srivastava, A. K., Janowski, M., and Lukomska, B. (2016). Extracellular vesicles in physiology, pathology, and therapy of the immune and central nervous system, with focus on extracellular vesicles derived from mesenchymal stem cells as therapeutic tools. *Front. Cell. Neurosci.* 10:109. doi: 10.3389/fncel.2016.00109

**Introduction** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=27199663>) | [CrossRef Full Text](#) (<https://doi.org/10.3389/fncel.2016.00109>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Extracellular+vesicles+in+physiology%2C+pathology%2C+and+therapy+of+the+immune+and+central+nervous+system%2C+with+focus+on+extracellular+vesicles+derived+from+mesenchymal+stem+cells+as+therapeutic+tools%2E&journal=Front%2E+Cell%2E+Neurosci%2E&author=Koniusz+S.+&author=Andrzejewska+A.&author=Muraca+M.&author=Srivastava+A.+K.&author=Janowski+M.&author=Lukomska+B.&publication\\_year=2016](http://scholar.google.com/scholar_lookup?&title=Extracellular+vesicles+in+physiology%2C+pathology%2C+and+therapy+of+the+immune+and+central+nervous+system%2C+with+focus+on+extracellular+vesicles+derived+from+mesenchymal+stem+cells+as+therapeutic+tools%2E&journal=Front%2E+Cell%2E+Neurosci%2E&author=Koniusz+S.+&author=Andrzejewska+A.&author=Muraca+M.&author=Srivastava+A.+K.&author=Janowski+M.&author=Lukomska+B.&publication_year=2016))

Krämer-Albers, E. M., and Hill, A. F. (2016). Extracellular vesicles: interneuronal shuttles of complex messages. *Curr. Opin. Neurobiol.* 39, 101–107. doi: 10.1016/j.conb.2016.04.016

**Response to Stroke** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=27183381>) | [CrossRef Full Text](#) (<https://doi.org/10.1016/j.conb.2016.04.016>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Extracellular+vesicles%3A+interneuronal+shuttles+of+complex+messages%2E&journal=Curr%2E+Opin%2E+Neurobiol%2E&author=Krämer-Albers+E.+M.&author=and+Hill+A.+F.&publication\\_year=2016&volume=39&pages=101-107](http://scholar.google.com/scholar_lookup?&title=Extracellular+vesicles%3A+interneuronal+shuttles+of+complex+messages%2E&journal=Curr%2E+Opin%2E+Neurobiol%2E&author=Krämer-Albers+E.+M.&author=and+Hill+A.+F.&publication_year=2016&volume=39&pages=101-107))

**In Response to**  
Kumar, A., Stoica, B. A., Loane, D. J., Yang, M., Abulwerdi, G., Khan, N., et al. (2017). Microglial-derived microparticles mediate neuroinflammation after traumatic brain injury. *J. Neuroinflammation* 14:47. doi: 10.1186/s12974-017-0819-4

**The Potential Use of**  
**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=28292310>) | [CrossRef Full Text](#) (<https://doi.org/10.1186/s12974-017-0819-4>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Microglial+derived+microparticles+mediate+neuroinflammation+after+traumatic+brain+injury%2E&journal=J%2E+Neuroinflammation&author=Kumar+A.&author=Stoica+B.+A.&author=Loane+D.+J.&author=Yang+M.&author=Abulwerdi+G.&author=Khan+N.&publication\\_year=2017](http://scholar.google.com/scholar_lookup?&title=Microglial+derived+microparticles+mediate+neuroinflammation+after+traumatic+brain+injury%2E&journal=J%2E+Neuroinflammation&author=Kumar+A.&author=Stoica+B.+A.&author=Loane+D.+J.&author=Yang+M.&author=Abulwerdi+G.&author=Khan+N.&publication_year=2017))

**Therapy**  
Lachenal, G., Pernet-Gallay, K., Chivet, M., Hemming, F. J., Belly, A., Bodon, G., et al. (2011). Release of exosomes from differentiated neurons and its regulation by synaptic glutamatergic activity. *Mol. Cell. Neurosci.* 46, 409–418. doi: 10.1016/j.mcn.2010.11.004

**Perspectives** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=21111824>) | [CrossRef Full Text](#) (<https://doi.org/10.1016/j.mcn.2010.11.004>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Release+of+exosomes+from+differentiated+neurons+and+its+regulation+by+synaptic+glutamatergic+activity%2E&journal=Mol%2E+Cell%2E+Neurosci%2E&author=Lachenal+G.&author=Pernet-Gallay+K.&author=Chivet+M.&author=Hemming+F.+J.&author=Belly+A.&author=Bodon+G.&publication\\_year=2011&volume=46&pages=409-418](http://scholar.google.com/scholar_lookup?&title=Release+of+exosomes+from+differentiated+neurons+and+its+regulation+by+synaptic+glutamatergic+activity%2E&journal=Mol%2E+Cell%2E+Neurosci%2E&author=Lachenal+G.&author=Pernet-Gallay+K.&author=Chivet+M.&author=Hemming+F.+J.&author=Belly+A.&author=Bodon+G.&publication_year=2011&volume=46&pages=409-418))

**Funding**  
Lai, C. P.-K., and Breakefield, X. O. (2012). Role of exosomes/microvesicles in the nervous system and use in emerging therapies. *Front. Physiol.* 3:228. doi: 10.3389/fphys.2012.00228

**Statement**  
**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=22754538>) | [CrossRef Full Text](#) (<https://doi.org/10.3389/fphys.2012.00228>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Role+of+exosomes%2Fmicrovesicles+in+the+nervous+system+and+use+in+emerging+therapies%2E&journal=Front%2E+Physiol%2E&author=Lai+C.+P.-K.&author=and+Breakefield+X.+O.&publication\\_year=2012](http://scholar.google.com/scholar_lookup?&title=Role+of+exosomes%2Fmicrovesicles+in+the+nervous+system+and+use+in+emerging+therapies%2E&journal=Front%2E+Physiol%2E&author=Lai+C.+P.-K.&author=and+Breakefield+X.+O.&publication_year=2012))

Lener, T., Gimona, M., Aigner, L., Börger, V., Buzas, E., Camussi, G., et al. (2015). Applying extracellular vesicles based therapeutics in clinical trials - an ISEV position paper. *J. Extracell. Vesicles* 4:30087. doi: 10.3402/jev.v4.30087

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=26725829>) | [CrossRef Full Text](#) (<https://doi.org/10.3402/jev.v4.30087>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Applying+extracellular+vesicles+based+therapeutics+in+clinical+trials+-+an+ISEV+position+paper%2E&journal=J%2E+Extracell%2E+Vesicles&author=Lener+T.&author=Gimona+M.&author=Aigner+L.&author=Börger+V.&author=Buzas+E.&author=Camussi+G.&publication\\_year=2015](http://scholar.google.com/scholar_lookup?&title=Applying+extracellular+vesicles+based+therapeutics+in+clinical+trials+-+an+ISEV+position+paper%2E&journal=J%2E+Extracell%2E+Vesicles&author=Lener+T.&author=Gimona+M.&author=Aigner+L.&author=Börger+V.&author=Buzas+E.&author=Camussi+G.&publication_year=2015))

Li, Y., Cheng, Q., Hu, G., Deng, T., Wang, Q., Zhou, J., et al. (2018). Extracellular vesicles in mesenchymal stromal cells: a novel therapeutic strategy for stroke. *Exp. Ther. Med.* 15, 4067–4079. doi: 10.3892/etm.2018.5993

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=29725359>) | [CrossRef Full Text](#) (<https://doi.org/10.3892/etm.2018.5993>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Extracellular+vesicles+in+mesenchymal+stromal+cells%3A+a+novel+therapeutic+strategy+for+stroke%2E&journal=Exp%2E+Ther%2E+Med%2E&author=Li+Y.&author=Cheng+Q.&author=Hu+G.&author=Deng+T.&author=Wang+Q.&author=Zhou+J.&publication\\_year=2018&volume=15&pages=4067-4079](http://scholar.google.com/scholar_lookup?&title=Extracellular+vesicles+in+mesenchymal+stromal+cells%3A+a+novel+therapeutic+strategy+for+stroke%2E&journal=Exp%2E+Ther%2E+Med%2E&author=Li+Y.&author=Cheng+Q.&author=Hu+G.&author=Deng+T.&author=Wang+Q.&author=Zhou+J.&publication_year=2018&volume=15&pages=4067-4079))

We use cookies on this website to improve your user experience. [Learn more](#) (<http://www.frontiersin.org/legal/list-of-cookies>)

CLOSE

- Ling, H., Fabbri, M., and Calin, G. A. (2013). MicroRNAs and other non-coding RNAs targets for anticancer drug development. *Nat. Rev. Drug Discov.* 12, 847–865. doi: 10.1038/nrd4140

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=24172333>) | [CrossRef Full Text](#) (<https://doi.org/10.1038/nrd4140>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=MicroRNAs+and+other+non-coding+RNAs+targets+for+anticancer+drug+development%2E&journal=Nat%2E+Rev%2E+Drug+Discov%2E&author=Ling+H.&author=Fabbri+M.&author=and+Calin+G.+A.&publication\\_year=2013&volume=12&pages=847-865](http://scholar.google.com/scholar_lookup?&title=MicroRNAs+and+other+non-coding+RNAs+targets+for+anticancer+drug+development%2E&journal=Nat%2E+Rev%2E+Drug+Discov%2E&author=Ling+H.&author=Fabbri+M.&author=and+Calin+G.+A.&publication_year=2013&volume=12&pages=847-865))

Liu, F. J., Lim, K. Y., Kaur, P., Sepramaniam, S., Armugam, A., Wong, P. T., et al. (2013). microRNAs involved in regulating spontaneous recovery in embolic stroke model. *PLoS One* 8:e66393. doi: 10.1371/journal.pone.0066393

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=23823624>) | [CrossRef Full Text](#) (<https://doi.org/10.1371/journal.pone.0066393>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=microRNAs+involved+in+regulating+spontaneous+recovery+in+embolic+stroke+model%2E&journal=PLoS+One&author=Liu+F.+J.&author=Lim+K.+Y.&author=Kaur+P.&author=Sepramaniam+S.&author=Armugam+A.&author=Wong+P.+T.&publication\\_year=2013](http://scholar.google.com/scholar_lookup?&title=microRNAs+involved+in+regulating+spontaneous+recovery+in+embolic+stroke+model%2E&journal=PLoS+One&author=Liu+F.+J.&author=Lim+K.+Y.&author=Kaur+P.&author=Sepramaniam+S.&author=Armugam+A.&author=Wong+P.+T.&publication_year=2013))

Llorente, A., van Deurs, B., and Sandvig, K. (2007). Cholesterol regulates prostasome release from secretory lysosomes in PC-3 human prostate cancer cells. *Eur. J. Cell Biol.* 86, 405–415. doi: 10.1016/j.ejcb.2007.05.001

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=17601634>) | [CrossRef Full Text](#) (<https://doi.org/10.1016/j.ejcb.2007.05.001>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Cholesterol+regulates+prostasome+release+from+secretory+lysosomes+in+PC-3+human+prostate+cancer+cells%2E&journal=Eur%2E+J%2E+Cell+Biol%2E&author=Llorente+A.&author=van+Deurs+B.&author=and+Sandvig+K.&publication\\_year=2007&volume=86&pages=405-415](http://scholar.google.com/scholar_lookup?&title=Cholesterol+regulates+prostasome+release+from+secretory+lysosomes+in+PC-3+human+prostate+cancer+cells%2E&journal=Eur%2E+J%2E+Cell+Biol%2E&author=Llorente+A.&author=van+Deurs+B.&author=and+Sandvig+K.&publication_year=2007&volume=86&pages=405-415))

Lopatina, T., Kalinina, N., Karagyaur, M., Stambolsky, D., Rubina, K., Revischin, A., et al. (2011). Adipose derived stem cells stimulate regeneration of peripheral nerves: BDNF secreted by these cells promotes nerve healing and axon growth *de novo*. *PLoS One* 6:e17899. doi: 10.1371/journal.pone.0017899

**Abstract**

**Introduction** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=21423756>) | [CrossRef Full Text](#) (<https://doi.org/10.1371/journal.pone.0017899>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Adipose+derived+stem+cells+stimulate+regeneration+of+peripheral+nerves%3A+BDNF+secreted+by+these+cells+promotes+nerve+healing+and+axon+growth+de+novo%2E&journal=PLoS+One&author=Lopatina+T.&author=Kalinina+N.&author=Karagyaur+M.&author=Stambolsky+D.&author=Rubina+T.C.&author=Revischin+A.&publication\\_year=2011](http://scholar.google.com/scholar_lookup?&title=Adipose+derived+stem+cells+stimulate+regeneration+of+peripheral+nerves%3A+BDNF+secreted+by+these+cells+promotes+nerve+healing+and+axon+growth+de+novo%2E&journal=PLoS+One&author=Lopatina+T.&author=Kalinina+N.&author=Karagyaur+M.&author=Stambolsky+D.&author=Rubina+T.C.&author=Revischin+A.&publication_year=2011))

**Tool**

Mayo, J. N., and Bearden, S. E. (2015). Driving the hypoxia inducible pathway in human pericytes promotes vascular density in an exosome dependent manner. *Microcirculation* 22, 711–723. doi: 10.1111/micc.12227

**Remodeling in Stroke**

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=26243428>) | [CrossRef Full Text](#) (<https://doi.org/10.1111/micc.12227>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Driving+the+hypoxia+inducible+pathway+in+human+pericytes+promotes+vascular+density+in+an+exosome+dependent+manner%2E&journal=Microcirculation&author=Mayo+J.+N.&author=and+Bearden+S.+E.&publication\\_year=2015&volume=22&pages=711-723](http://scholar.google.com/scholar_lookup?&title=Driving+the+hypoxia+inducible+pathway+in+human+pericytes+promotes+vascular+density+in+an+exosome+dependent+manner%2E&journal=Microcirculation&author=Mayo+J.+N.&author=and+Bearden+S.+E.&publication_year=2015&volume=22&pages=711-723))

**Derived Exosomes**

Mosser, D. M., and Zhang, X. (2008). Interleukin-10: new perspectives on an old cytokine. *Immunol. Rev.* 226, 205–218. doi: 10.1111/j.1600-065X.2008.00706.x

**Stroke**

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=19161426>) | [CrossRef Full Text](#) (<https://doi.org/10.1111/j.1600-065X.2008.00706.x>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Interleukin-10%3A+new+perspectives+on+an+old+cytokine%2E&journal=Immunol%2E++Rev%2E&author=Mosser+D.+M.&author=and+Zhang+X.&publication\\_year=2008&volume=226&pages=205-218](http://scholar.google.com/scholar_lookup?&title=Interleukin-10%3A+new+perspectives+on+an+old+cytokine%2E&journal=Immunol%2E++Rev%2E&author=Mosser+D.+M.&author=and+Zhang+X.&publication_year=2008&volume=226&pages=205-218))

**Exosomes for Stroke**

Nakagomi, T., Kubo, S., Nakano-Doi, A., Sakuma, R., Lu, S., Narita, A., et al. (2015). Brain vascular pericytes following ischemia have multipotential stem cell activity to differentiate into neural and vascular lineage cells. *Stem Cells* 33, 1962–1974. doi: 10.1002/stem.1977

**Conclusion and Author Contributions**

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=25694098>) | [CrossRef Full Text](#) (<https://doi.org/10.1002/stem.1977>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Conclusion+and+Author+Contributions&author=Nakagomi+T.&author=Kubo+S.&author=Nakano-Doi+A.&author=Sakuma+R.&author=Lu+S.&author=Narita+A.&publication\\_year=2015&volume=33&pages=1962-1974](http://scholar.google.com/scholar_lookup?&title=Conclusion+and+Author+Contributions&author=Nakagomi+T.&author=Kubo+S.&author=Nakano-Doi+A.&author=Sakuma+R.&author=Lu+S.&author=Narita+A.&publication_year=2015&volume=33&pages=1962-1974))

**Funding**

Nakajima, M., Nito, C., Sowa, K., Suda, S., Nishiyama, Y., Nakamura-Takahashi, A., et al. (2017). Mesenchymal stem cells overexpressing interleukin-10 promote neuroprotection in experimental acute ischemic stroke. *Mol. Ther. Methods Clin. Dev.* 6, 102–111. doi: 10.1016/j.omtm.2017.06.005

**Conflict of Interest**

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=28725658>) | [CrossRef Full Text](#) (<https://doi.org/10.1016/j.omtm.2017.06.005>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Mesenchymal+stem+cells+overexpressing+interleukin-10+promote+neuroprotection+in+experimental+acute+ischemic+stroke%2E&journal=Mol%2E+Ther%2E+Methods+Clin%2E+Dev%2E&author=Nakajima+M.&author=Nito+C.&author=Sowa+K.&author=Suda+S.&author=Nishiyama+Y.&author=Nakamura-Takahashi+A.&publication\\_year=2017&volume=6&pages=102-111](http://scholar.google.com/scholar_lookup?&title=Mesenchymal+stem+cells+overexpressing+interleukin-10+promote+neuroprotection+in+experimental+acute+ischemic+stroke%2E&journal=Mol%2E+Ther%2E+Methods+Clin%2E+Dev%2E&author=Nakajima+M.&author=Nito+C.&author=Sowa+K.&author=Suda+S.&author=Nishiyama+Y.&author=Nakamura-Takahashi+A.&publication_year=2017&volume=6&pages=102-111))

Norden, D. M., Fenn, A. M., Dugan, A., and Godbout, J. P. (2014). TGF beta produced by IL-10 redirected astrocytes attenuates microglial activation. *Glia* 62, 881–895. doi: 10.1002/glia.22647

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=24616125>) | [CrossRef Full Text](#) (<https://doi.org/10.1002/glia.22647>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=TGF+beta+produced+by+IL-10+redirected+astrocytes+attenuates+microglial+activation%2E&journal=Glia&author=Norden+D.+M.&author=Fenn+A.+M.&author=Dugan+A.&author=and+Godbout+J.+P.&publication\\_year=2014&volume=62&pages=881-895](http://scholar.google.com/scholar_lookup?&title=TGF+beta+produced+by+IL-10+redirected+astrocytes+attenuates+microglial+activation%2E&journal=Glia&author=Norden+D.+M.&author=Fenn+A.+M.&author=Dugan+A.&author=and+Godbout+J.+P.&publication_year=2014&volume=62&pages=881-895))

Otero-Ortega, L., Gomez de Frutos, M. C., Laso-Garcia, F., Rodriguez-Frutos, B., Medina-Gutierrez, E., Lopez, J. A., et al. (2017). Exosomes promote restoration after an experimental animal model of intracerebral haemorrhage. *J. Cereb. Blood Flow Metab.* 38, 767–779. doi: 10.1177/0271678X17708917

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=28524762>) | [CrossRef Full Text](#) (<https://doi.org/10.1177/0271678X17708917>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Exosomes+promote+restoration+after+an+experimental+animal+model+of+intracerebral+haemorrhage%2E&journal=J%2E+Cereb%2E+Blood+Flow+Metab%2E&author=Otero-Ortega+L.&author=Gomez+de+Frutos+M.+C.&author=Laso-Garcia+F.&author=Rodriguez-Frutos+B.&author=Medina-Gutierrez+E.&author=Lopez+J.+A.+A.&publication\\_year=2017&volume=38&pages=767-779](http://scholar.google.com/scholar_lookup?&title=Exosomes+promote+restoration+after+an+experimental+animal+model+of+intracerebral+haemorrhage%2E&journal=J%2E+Cereb%2E+Blood+Flow+Metab%2E&author=Otero-Ortega+L.&author=Gomez+de+Frutos+M.+C.&author=Laso-Garcia+F.&author=Rodriguez-Frutos+B.&author=Medina-Gutierrez+E.&author=Lopez+J.+A.+A.&publication_year=2017&volume=38&pages=767-779))

Popa-Wagner, A., Buga, A. M., and Kokaia, Z. (2011). Perturbed cellular response to brain injury during aging. *Aging Res. Rev.* 10, 71–79. doi: 10.1016/j.arr.2009.10.008

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=19900590>) | [CrossRef Full Text](#) (<https://doi.org/10.1016/j.arr.2009.10.008>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Perturbed+cellular+response+to+brain+injury+during+aging%2E&journal=Aging+Res%2E++Rev%2E&author=Popa-Wagner+A.&author=Buga+A.+M.&author=and+Kokaia+Z.&publication\\_year=2011&volume=10&pages=71-79](http://scholar.google.com/scholar_lookup?&title=Perturbed+cellular+response+to+brain+injury+during+aging%2E&journal=Aging+Res%2E++Rev%2E&author=Popa-Wagner+A.&author=Buga+A.+M.&author=and+Kokaia+Z.&publication_year=2011&volume=10&pages=71-79))

Popa-Wagner, A., Carmichael, S. T., Kokaia, Z., and Walker, L. C. (2007). The response of the aged brain to stroke: too much, too soon? *Curr. Neurovasc. Res.* 4, 216–277. doi: 10.2174/156720207781387213

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=17691975>) | [CrossRef Full Text](#) (<https://doi.org/10.2174/156720207781387213>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=The+response+of+the+aged+brain+to+stroke%3A+too+much%2C+too+soon%2B4&journal=Curr%2E+Neurovasc%2E++Res%2E&author=Popa-Wagner+A.&author=Carmichael+S.+T.&author=Kokaia+Z.&author=and+Walker+L.+C.&publication\\_year=2007&volume=4&pages=216-277](http://scholar.google.com/scholar_lookup?&title=The+response+of+the+aged+brain+to+stroke%3A+too+much%2C+too+soon%2B4&journal=Curr%2E+Neurovasc%2E++Res%2E&author=Popa-Wagner+A.&author=Carmichael+S.+T.&author=Kokaia+Z.&author=and+Walker+L.+C.&publication_year=2007&volume=4&pages=216-277))

Popa-Wagner, A., Dinca, I., Yalikun, S., Walker, L., Kroemer, H., and Kessler, C. (2006). Accelerated delimitation of the infarct zone by capillary-derived nestin-positive cells in aged rats. *Curr. Neurovasc. Res.* 3, 3–13. doi: 10.2174/15672020675541732

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=16472121>) | [CrossRef Full Text](#) (<https://doi.org/10.2174/15672020675541732>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Accelerated+delimitation+of+the+infarct+zone+by+capillary-derived+nestin-positive+cells+in+aged+rats%2E&journal=Curr%2E++Neurovasc%2E++Res%2E&author=Popa-Wagner+A.&author=Dinca+I.&author=Yalikun+S.&author=Walker+L.&author=Kroemer+H.&author=and+Kessler+C.&publication\\_year=2006&volume=3&pages=3-13](http://scholar.google.com/scholar_lookup?&title=Accelerated+delimitation+of+the+infarct+zone+by+capillary-derived+nestin-positive+cells+in+aged+rats%2E&journal=Curr%2E++Neurovasc%2E++Res%2E&author=Popa-Wagner+A.&author=Dinca+I.&author=Yalikun+S.&author=Walker+L.&author=Kroemer+H.&author=and+Kessler+C.&publication_year=2006&volume=3&pages=3-13))

**Abstract**  
Popa-Wagner, A., Glavan, D. G., Olaru, A., Olaru, D. G., Margaritescu, O., Tica, O., et al. (2018). Present status and future challenges of new therapeutic targets in preclinical models of stroke in aged animals with/without comorbidities. *Int. J. Mol. Sci.* 19:E356. doi: 10.3390/ijms19020356  
**INTRODUCTION**

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=29370078>) | [CrossRef Full Text](#) (<https://doi.org/10.3390/ijms19020356>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Present+status+and+future+challenges+of+new+therapeutic+targets+in+preclinical+models+of+stroke+in+aged+animals+with%2Fwithout+co-morbidities%2E&journal=Int%2E+J%2E+Mol%2E+Sci%2E&author=Popa-Wagner+A.&author=Glavan+D.+G.&author=Olaru+A.&author=Olaru+D.+G.&author=Margaritescu+O.&author=Tica+O.&publication\\_year=2018](http://scholar.google.com/scholar_lookup?&title=Present+status+and+future+challenges+of+new+therapeutic+targets+in+preclinical+models+of+stroke+in+aged+animals+with%2Fwithout+co-morbidities%2E&journal=Int%2E+J%2E+Mol%2E+Sci%2E&author=Popa-Wagner+A.&author=Glavan+D.+G.&author=Olaru+A.&author=Olaru+D.+G.&author=Margaritescu+O.&author=Tica+O.&publication_year=2018))

**Proteomic Analysis of Microglia-Derived Exosomes**  
Potolicchio, I., Carven, G. J., Xu, X., Stipp, C., Riese, R. J., Stern, L. J., et al. (2005). Proteomic analysis of microglia-derived exosomes: metabolic role of the aminopeptidase CD13 in neuropeptide catabolism. *J. Immunol.* 175, 2237–2243. doi: 10.4049/jimmunol.175.4.2237

**Response to Stroke**  
(<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=16081791>) | [CrossRef Full Text](#) (<https://doi.org/10.4049/jimmunol.175.4.2237>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Proteomic+analysis+of+microglia-derived+exosomes%3A+metabolic+role+of+the+aminopeptidase+CD13+in+neuropeptide+catabolism%2E&journal=J%2E+Immunol%2E&author=Potolicchio,I.&author=Carven+G.+J.&author=Xu+X.&author=Stipp+C.&author=Riese+R.+J.&author=Stern+L.+J.&publication\\_year=2005&volume=175&pages=2237-2243](http://scholar.google.com/scholar_lookup?&title=Proteomic+analysis+of+microglia-derived+exosomes%3A+metabolic+role+of+the+aminopeptidase+CD13+in+neuropeptide+catabolism%2E&journal=J%2E+Immunol%2E&author=Potolicchio,I.&author=Carven+G.+J.&author=Xu+X.&author=Stipp+C.&author=Riese+R.+J.&author=Stern+L.+J.&publication_year=2005&volume=175&pages=2237-2243))

**Stroke**  
Record, M., Poirot, M., and Silvente-Poirot, S. (2014). Emerging concepts on the role of exosomes in lipid metabolic diseases. *Biochimie* 96, 67–74. doi: 10.1016/j.biochi.2013.06.016

**Mesenchymal Stem Cells and Their Emerging Role in Stroke**  
(<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=23827857>) | [CrossRef Full Text](#) (<https://doi.org/10.1016/j.jbiochi.2013.06.016>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Cells+and+Their+Emerging+Role+in+Stroke%2E&journal=Biochimie&author=Record+M.&author=Poirot+M.+Silvente+Poirot+S.&publication\\_year=2014&volume=96&pages=67-74](http://scholar.google.com/scholar_lookup?&title=Cells+and+Their+Emerging+Role+in+Stroke%2E&journal=Biochimie&author=Record+M.&author=Poirot+M.+Silvente+Poirot+S.&publication_year=2014&volume=96&pages=67-74))

**Post-Acute Care and Remodeling**  
Reitmeir, R., Kilic, E., Kilic, U., Bacigaluppi, M., ElAli, A., Salani, G., et al. (2011). Post-acute delivery of erythropoietin induces stroke recovery by promoting perilesional tissue remodelling and contralesional pyramidal tract plasticity. *Brain* 134, 84–99. doi: 10.1093/brain/awq344

**Post-Acute Care and Remodeling**  
(<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=21186263>) | [CrossRef Full Text](#) (<https://doi.org/10.1093/brain/awq344>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Post-acute+delivery+of+erythropoietin+induces+stroke+recovery+by+promoting+perilesional+tissue+remodelling+and+contralesional+pyramidal+tract+plasticity%2E&journal=Brain&author=Reitmeir+R.&author=Kilic+E.&author=Kilic+U.&author=Bacigaluppi+M.&author=ElAli+A.&author=Salani+G.&publication\\_year=2011&volume=134&pages=84-99](http://scholar.google.com/scholar_lookup?&title=Post-acute+delivery+of+erythropoietin+induces+stroke+recovery+by+promoting+perilesional+tissue+remodelling+and+contralesional+pyramidal+tract+plasticity%2E&journal=Brain&author=Reitmeir+R.&author=Kilic+E.&author=Kilic+U.&author=Bacigaluppi+M.&author=ElAli+A.&author=Salani+G.&publication_year=2011&volume=134&pages=84-99))

**Conflict of Interest Statement**  
Rosell, A., and Lo, E. H. (2008). Multiphasic roles for matrix metalloproteinases after stroke. *Curr. Opin. Pharmacol.* 8, 82–89. doi: 10.1016/j.coph.2007.12.001

**Footnotes**  
(<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=18226583>) | [CrossRef Full Text](#) (<https://doi.org/10.1016/j.coph.2007.12.001>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Multiphasic+roles+for+matrix+metalloproteinases+after+stroke%2E&journal=Curr%2E++Opin%2E++Pharmacol%2E&author=Rosell+A.&author=Lo+E.+H.&publication\\_year=2008&volume=8&pages=82-89](http://scholar.google.com/scholar_lookup?&title=Multiphasic+roles+for+matrix+metalloproteinases+after+stroke%2E&journal=Curr%2E++Opin%2E++Pharmacol%2E&author=Rosell+A.&author=Lo+E.+H.&publication_year=2008&volume=8&pages=82-89))

**MicroRNAs and Exosomes**  
Salido-Guadarrama, I., Romero-Cordoba, S., Peralta-Zaragoza, O., Hidalgo-Miranda, A., and Rodriguez-Dorantes, M. (2014). MicroRNAs transported by exosomes in body fluids as mediators of intercellular communication in cancer. *Onco Targets Ther.* 7, 1327–1338. doi: 10.2147/OTT.S61562

**MicroRNAs and Exosomes**  
(<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=25092989>) | [CrossRef Full Text](#) (<https://doi.org/10.2147/OTT.S61562>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=MicroRNAs+transported+by+exosomes+in+body+fluids+as+mediators+of+intercellular+communication+in+cancer%2E&journal=Oncotargets+Ther%2E&author=Salido-Guadarrama+I.&author=Romero-Cordoba+S.&author=Peralta-Zaragoza+O.&author=Hidalgo-Miranda+A.&author=Rodriguez-Dorantes+M.&publication\\_year=2014&volume=7&pages=1327-1338](http://scholar.google.com/scholar_lookup?&title=MicroRNAs+transported+by+exosomes+in+body+fluids+as+mediators+of+intercellular+communication+in+cancer%2E&journal=Oncotargets+Ther%2E&author=Salido-Guadarrama+I.&author=Romero-Cordoba+S.&author=Peralta-Zaragoza+O.&author=Hidalgo-Miranda+A.&author=Rodriguez-Dorantes+M.&publication_year=2014&volume=7&pages=1327-1338))

Savitz, S. I., and Fisher, M. (2007). Future of neuroprotection for acute stroke: in the aftermath of the SAINT trials. *Ann. Neurol.* 61, 396–402. doi: 10.1002/ana.21127

**Future of Neuroprotection**  
(<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=17420989>) | [CrossRef Full Text](#) (<https://doi.org/10.1002/ana.21127>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Future+of+neuroprotection+for+acute+stroke%3A+in+the+aftermath+of+the+SAINT+trials%2E&journal=Ann%2E+Neurol%2E&author=Savitz+S.+I.&author=Fisher+M.&publication\\_year=2007&volume=61&pages=396-402](http://scholar.google.com/scholar_lookup?&title=Future+of+neuroprotection+for+acute+stroke%3A+in+the+aftermath+of+the+SAINT+trials%2E&journal=Ann%2E+Neurol%2E&author=Savitz+S.+I.&author=Fisher+M.&publication_year=2007&volume=61&pages=396-402))

We use cookies on this website to improve your user experience. [Learn more](#) (<http://www.frontiersin.org/legal/list-of-cookies>)

CLOSE

Skotland, T., Sandvig, K., and Llorente, A. (2017). Lipids in exosomes: current knowledge and the way forward. *Prog. Lipid Res.* 66, 30–41. doi: 10.1016/j.plipres.2017.03.001

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=28342835>) | [CrossRef Full Text](#) (<https://doi.org/10.1016/j.plipres.2017.03.001>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Lipids+in+exosomes%3A+current+knowledge+and+the+way+forward%2E&journal=Prog%2E++Lipid+Res%2E&author=Skotland+T.&author=S+andvig+K.&author=and+Llorente+A.&publication\\_year=2017&volume=66&pages=30-41](http://scholar.google.com/scholar_lookup?&title=Lipids+in+exosomes%3A+current+knowledge+and+the+way+forward%2E&journal=Prog%2E++Lipid+Res%2E&author=Skotland+T.&author=S+andvig+K.&author=and+Llorente+A.&publication_year=2017&volume=66&pages=30-41))

Strauss, K., Goebel, C., Runz, H., Möbius, W., Weiss, S., Feussner, I., et al. (2010). Exosome secretion ameliorates lysosomal storage of cholesterol in niemann-pick type c disease. *J. Biol. Chem.* 285, 26279–26288. doi: 10.1074/jbc.M110.134775

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=20554533>) | [CrossRef Full Text](#) (<https://doi.org/10.1074/jbc.M110.134775>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Exosome+secretion+ameliorates+lysosomal+storage+of+cholesterol+in+niemann-pick+type+c+disease%2E&journal=J%2E++Biol%2E++Chem%2E&author=Strauss+K.&author=Goebel+C.&author=Runz+H.&author=Möbius+W.&author=Weiss+S.&author=Feussner+I.&publication\\_year=2010&volume=285&pages=26279-26288](http://scholar.google.com/scholar_lookup?&title=Exosome+secretion+ameliorates+lysosomal+storage+of+cholesterol+in+niemann-pick+type+c+disease%2E&journal=J%2E++Biol%2E++Chem%2E&author=Strauss+K.&author=Goebel+C.&author=Runz+H.&author=Möbius+W.&author=Weiss+S.&author=Feussner+I.&publication_year=2010&volume=285&pages=26279-26288))

Taylor, A. R., Robinson, M. B., Gifondorwa, D. J., Tytell, M., and Milligan, C. E. (2007). Regulation of heat shock protein 70 release in astrocytes: role of signaling kinases. *Dev. Neurobiol.* 67, 1815–1829. doi: 10.1002/dneu.20559

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=17701989>) | [CrossRef Full Text](#) (<https://doi.org/10.1002/dneu.20559>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Regulation+of+heat+shock+protein+70+release+in+astrocytes%3A+role+of+signaling+kinases%2E&journal=Dev%2E+Neurobiol%2E&author=T+TABLE+OF+CONTENTS+Robinson+M.+B.&author=Gifondorwa+D.+J.&author=Tytell+M.&author=and+Milligan+C.+E.&publication\\_year=2007&volume=67&pages=1815-1829](http://scholar.google.com/scholar_lookup?&title=Regulation+of+heat+shock+protein+70+release+in+astrocytes%3A+role+of+signaling+kinases%2E&journal=Dev%2E+Neurobiol%2E&author=T+TABLE+OF+CONTENTS+Robinson+M.+B.&author=Gifondorwa+D.+J.&author=Tytell+M.&author=and+Milligan+C.+E.&publication_year=2007&volume=67&pages=1815-1829))

**Abstract**  
Théry, C., Boussac, M., Véron, P., Ricciardi-Castagnoli, P., Raposo, G., Garin, J., et al. (2001). Proteomic analysis of dendritic cell-derived exosomes: a secreted subcellular compartment distinct from apoptotic vesicles. *J. Immunol.* 166, 7309–7318. doi: 10.4049/jimmunol.166.12.7309  
**Introduction**

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=11390481>) | [CrossRef Full Text](#) (<https://doi.org/10.4049/jimmunol.166.12.7309>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Proteomic+analysis+of+dendritic+cell-derived+exosomes+as+a+secreted+subcellular+compartment+distinct+from+apoptotic+vesicles%2E&journal=J%2E+Immunol%2E&author=Théry+C.+&author=Boussac+M.+&author=Véron+P.+&author=Ricciardi-Castagnoli+P.+&author=Raposo+G.+&author=Garin+J.+&author=and+Dignat-George+F.+&publication\\_year=2001&volume=166&pages=7309-7318](http://scholar.google.com/scholar_lookup?&title=Proteomic+analysis+of+dendritic+cell-derived+exosomes+as+a+secreted+subcellular+compartment+distinct+from+apoptotic+vesicles%2E&journal=J%2E+Immunol%2E&author=Théry+C.+&author=Boussac+M.+&author=Véron+P.+&author=Ricciardi-Castagnoli+P.+&author=Raposo+G.+&author=Garin+J.+&author=and+Dignat-George+F.+&publication_year=2001&volume=166&pages=7309-7318))  
**Neurovascular Unit**  
Todorova, D., Simoncini, S., Lacroix, R., Sabatier, F., and Dignat-George, F. (2017). Extracellular vesicles in angiogenesis. *Circ. Res.* 120, 1658–1673. doi: 10.1161/CIRCRESAHA.117.309681  
**Response to Stroke**

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=28495996>) | [CrossRef Full Text](#) (<https://doi.org/10.1161/CIRCRESAHA.117.309681>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Derived+Exosomes+in+angiogenesis%2E&journal=Circ%2E+Res%2E&author=Todorova+D.+&author=Simoncini+S.+&author=Lacroix+R.+&author=Sabatier+F.+&author=and+Dignat-George+F.+&publication\\_year=2017&volume=120&pages=1658-1673](http://scholar.google.com/scholar_lookup?&title=Derived+Exosomes+in+angiogenesis%2E&journal=Circ%2E+Res%2E&author=Todorova+D.+&author=Simoncini+S.+&author=Lacroix+R.+&author=Sabatier+F.+&author=and+Dignat-George+F.+&publication_year=2017&volume=120&pages=1658-1673))

**Stroke**  
Tsenkina, Y., Ricard, J., Runko, E., Quiala-Acosta, M. M., Mier, J., and Liebl, D. J. (2015). EphB3 receptors function as dependence receptors to mediate oligodendrocyte cell death following contusive spinal cord injury. *Cell Death Dis.* 6:e1922. doi: 10.1038/cddis.2015.262

**Mesenchymal Stem Cells and Their Derivatives**  
**Stroke**  
Tsenkina, Y., Ricard, J., Runko, E., Quiala-Acosta, M. M., Mier, J., and Liebl, D. J. (2015). EphB3 receptors function as dependence receptors to mediate oligodendrocyte cell death following contusive spinal cord injury%2E&journal=Cell+Death+Dis%2E&author=Tsenkina+Y.+&author=Ricard+J.+&author=Runko+E.+&author=Quiala-Acosta+M.+M.+&author=Mier+J.+&author=and+Liebl+D.+&publication\_year=2015  
**Conclusion and Perspectives**  
Valadi, H., Ekström, K., Bossios, A., Sjöstrand, M., Lee, J. J., and Lötvall, J. O. (2007). Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat. Cell Biol.* 9, 654–659. doi: 10.1038/ncb1596

**Author Contributions**  
**Funding**  
**Statement**  
Van Meer, G., Voelker, D. R., and Feigenson, G. W. (2008). Membrane lipids: where they are and how they behave. *Nat. Rev. Mol. Cell Biol.* 9, 112–124. doi: 10.1038/nrm2330  
**Footnotes**

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=18216768>) | [CrossRef Full Text](#) (<https://doi.org/10.1038/nrm2330>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Membrane+lipids%3A+where+they+are+and+how+they+behave%2E&journal=Nat%2E+Rev%2E+Mol%2E+Cell+Biol%2E&author=Van+Meer+G.+&author=Voelker+D.+R.+&author=and+Feigenson+G.+W.+&publication\\_year=2008&volume=9&pages=112-124](http://scholar.google.com/scholar_lookup?&title=Membrane+lipids%3A+where+they+are+and+how+they+behave%2E&journal=Nat%2E+Rev%2E+Mol%2E+Cell+Biol%2E&author=Van+Meer+G.+&author=Voelker+D.+R.+&author=and+Feigenson+G.+W.+&publication_year=2008&volume=9&pages=112-124))

Verderio, C., Muzio, L., Turola, E., Bergami, A., Novellino, L., Ruffini, F., et al. (2012). Myeloid microvesicles are a marker and therapeutic target for neuroinflammation. *Ann. Neurol.* 72, 610–624. doi: 10.1002/ana.23627

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=23109155>) | [CrossRef Full Text](#) (<https://doi.org/10.1002/ana.23627>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Myeloid+microvesicles+are+a+marker+and+therapeutic+target+for+neuroinflammation%2E&journal=Ann%2E+Neurol%2E&author=Verderio+C.+&author=Muzio+L.+&author=Turola+E.+&author=Bergami+A.+&author=Novellino+L.+&author=Ruffini+F.+&publication\\_year=2012&volume=72&pages=610-624](http://scholar.google.com/scholar_lookup?&title=Myeloid+microvesicles+are+a+marker+and+therapeutic+target+for+neuroinflammation%2E&journal=Ann%2E+Neurol%2E&author=Verderio+C.+&author=Muzio+L.+&author=Turola+E.+&author=Bergami+A.+&author=Novellino+L.+&author=Ruffini+F.+&publication_year=2012&volume=72&pages=610-624))

Von Bartheld, C. S., and Altick, A. L. (2011). Multivesicular bodies in neurons: distribution, protein content, and trafficking functions. *Prog. Neurobiol.* 93, 313–340. doi: 10.1016/j.pneurobio.2011.01.003

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=21216273>) | [CrossRef Full Text](#) (<https://doi.org/10.1016/j.pneurobio.2011.01.003>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Multivesicular+bodies+in+neurons%3A+distribution%2C+protein+content%2C+and+trafficking+functions%2E&journal=Prog%2E++Neurobiol%2E&author=Von+Bartheld+C.+S.+&author=Altick+A.+L.+&publication\\_year=2011&volume=93&pages=313-340](http://scholar.google.com/scholar_lookup?&title=Multivesicular+bodies+in+neurons%3A+distribution%2C+protein+content%2C+and+trafficking+functions%2E&journal=Prog%2E++Neurobiol%2E&author=Von+Bartheld+C.+S.+&author=Altick+A.+L.+&publication_year=2011&volume=93&pages=313-340))

Xin, H., Katakowski, M., Wang, F., Qian, J.-Y., Shuang Liu, X., Ali, M. M., et al. (2017a). MiR-17-92 cluster in exosomes enhance neuroplasticity and functional recovery after stroke in rats. *Stroke* 48, 747–753. doi: 10.1161/STROKEAHA.116.015204

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=28232590>) | [CrossRef Full Text](#) (<https://doi.org/10.1161/STROKEAHA.116.015204>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=MiR-17-92+cluster+in+exosomes+enhance+neuroplasticity+and+functional+recovery+after+stroke+in+rats%2E&journal=Stroke&author=Xin+H.&author=Katakowsk+M.&author=Wang+F.&author=Qian+J.-Y.&author=Shuang+Liu+X.&author=Ali+M.+M.&publication\\_year=2017a&volume=48&pages=747-753](http://scholar.google.com/scholar_lookup?&title=MiR-17-92+cluster+in+exosomes+enhance+neuroplasticity+and+functional+recovery+after+stroke+in+rats%2E&journal=Stroke&author=Xin+H.&author=Katakowsk+M.&author=Wang+F.&author=Qian+J.-Y.&author=Shuang+Liu+X.&author=Ali+M.+M.&publication_year=2017a&volume=48&pages=747-753))

Xin, H., Wang, F., Li, Y., Lu, Q.-E., Cheung, W. L., Zhang, Y., et al. (2017b). Secondary release of exosomes from astrocytes contributes to the increase in neural plasticity and improvement of functional recovery after stroke in rats treated with exosomes harvested from MicroRNA 133b-overexpressing multipotent mesenchymal stromal cells. *Cell Trans.* 26, 243–257. doi: 10.3727/096368916X693031

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=27677799>) | [CrossRef Full Text](#) (<https://doi.org/10.3727/096368916X693031>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Secondary+release+of+exosomes+from+astrocytes+contributes+to+the+increase+in+neural+plasticity+and+improvement+of+functional+recovery+after+stroke+in+rats+treated+with+exosomes+harvested+from+MicroRNA+133b-overexpressing+multipotent+mesenchymal+stromal+cells%2E&journal=Cell+Trans%2E&author=Xin+H.&author=Wang+F.&author=Li+Y.&author=Lu+Q.-E.&author=Cheung+W.+L.&author=Zhang+Y.&publication\\_year=2017b&volume=26&pages=243-257](http://scholar.google.com/scholar_lookup?&title=Secondary+release+of+exosomes+from+astrocytes+contributes+to+the+increase+in+neural+plasticity+and+improvement+of+functional+recovery+after+stroke+in+rats+treated+with+exosomes+harvested+from+MicroRNA+133b-overexpressing+multipotent+mesenchymal+stromal+cells%2E&journal=Cell+Trans%2E&author=Xin+H.&author=Wang+F.&author=Li+Y.&author=Lu+Q.-E.&author=Cheung+W.+L.&author=Zhang+Y.&publication_year=2017b&volume=26&pages=243-257))

Xin, H., Li, Y., Cui, Y., Yang, J. J., Zhang, Z. G., and Chopp, M. (2013). Systemic administration of exosomes released from mesenchymal stromal cells promote functional recovery and neurovascular plasticity after stroke in rats. *J. Cereb. Blood Flow Metab.* 33, 1711–1715. doi: 10.1038/jcbfm.2013.152

**TABLE OF CONTENTS** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=23963371>) | [CrossRef Full Text](#) (<https://doi.org/10.1038/jcbfm.2013.152>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Systemic+administration+of+exosomes+released+from+mesenchymal+stromal+cells+promote+functional+recovery+and+neurovascular+plasticity+after+stroke+in+rats%2E&journal=J%2E+Cereb%2E+Blood+Flow+Metab%2E&author=Xin+H.&author=Li+Y.&author=Cui+Y.&author=Yang+J.+J.&author=Zhang+Z.+G.&author=and+Chopp+M.&publication\\_year=2013&volume=33&pages=1711-1715](http://scholar.google.com/scholar_lookup?&title=Systemic+administration+of+exosomes+released+from+mesenchymal+stromal+cells+promote+functional+recovery+and+neurovascular+plasticity+after+stroke+in+rats%2E&journal=J%2E+Cereb%2E+Blood+Flow+Metab%2E&author=Xin+H.&author=Li+Y.&author=Cui+Y.&author=Yang+J.+J.&author=Zhang+Z.+G.&author=and+Chopp+M.&publication_year=2013&volume=33&pages=1711-1715))

**Yang S, Jin-a Ha Zhu, Y., Wan, Y., Opoku, E. N., Zhu, L., et al. (2017). Diverse functions and mechanisms of pericytes in ischemic stroke. *Curr. Neuropharmacol.* 15, 892–905. doi: 10.2174/1570159X15666170112170226**

**Communication** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=28088914>) | [CrossRef Full Text](#) (<https://doi.org/10.2174/1570159X15666170112170226>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Diverse+functions+and+mechanisms+of+pericytes+in+ischemic+stroke%2E&journal=Curr%2E+Neuropharmacol%2E&author=Yang+S.&author=Neurovascular+Unit%2E+and+Zhu+Y.&author=Zhu+Y.&author=Opoku+E.+N.&author=Zhu+L.&publication\\_year=2017&volume=15&pages=892-905](http://scholar.google.com/scholar_lookup?&title=Diverse+functions+and+mechanisms+of+pericytes+in+ischemic+stroke%2E&journal=Curr%2E+Neuropharmacol%2E&author=Yang+S.&author=Neurovascular+Unit%2E+and+Zhu+Y.&author=Zhu+Y.&author=Opoku+E.+N.&author=Zhu+L.&publication_year=2017&volume=15&pages=892-905))

Remodeling in **Zhang, Z. G., and Chopp, M. (2016). Exosomes in stroke pathogenesis and therapy. *J. Clin. Invest.* 126, 1190–1197. doi: 10.1172/JCI81133**

**PubMed Abstract** (<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd>ShowDetailView&TermToSearch=27035810>) | [CrossRef Full Text](#) (<https://doi.org/10.1172/JCI81133>) | [Google Scholar](#) ([http://scholar.google.com/scholar\\_lookup?&title=Exosomes+in+stroke+pathogenesis+and+therapy%2E&journal=J%2E+Clin%2E+Invest%2E&author=Zhang+Z.+G.&author=and+Chopp+M.&publication\\_year=2016&volume=126&pages=1190-1197](http://scholar.google.com/scholar_lookup?&title=Exosomes+in+stroke+pathogenesis+and+therapy%2E&journal=J%2E+Clin%2E+Invest%2E&author=Zhang+Z.+G.&author=and+Chopp+M.&publication_year=2016&volume=126&pages=1190-1197))

**Stroke** **Zlokovic, B. V. (2008). The blood–brain barrier in health and chronic neurodegenerative disorders. *Neuron* 57, 178–201. doi: 10.1016/j.neuro.2008.01.003**

**Mesenchymal Stem Cells and Their Contribution to Stroke** **Zagrean A-M, Hermann DM, Opris I, Zagrean L and Popa-Wagner A (2018) Multicellular Crosstalk Between Exosomes and the Neurovascular Unit After Cerebral Ischemia. Therapeutic Implications. *Front. Neurosci.* 12:811. doi: 10.3389/fnins.2018.00811**

## Conclusion and Perspectives

**Keywords:** exosome, nanovesicles, neurovascular unit, blood-brain barrier, miRNA, stroke, mesenchymal stem cells

**Author Contributions**

**Citation:** Zagrean A-M, Hermann DM, Opris I, Zagrean L and Popa-Wagner A (2018) Multicellular Crosstalk Between Exosomes and the Neurovascular Unit After Cerebral Ischemia. Therapeutic Implications. *Front. Neurosci.* 12:811. doi: 10.3389/fnins.2018.00811

**Funding**

**Received:** 02 July 2018; **Accepted:** 17 October 2018;

**Published:** 06 November 2018.

**Conflict of Interest**

**Statement**

**Hari S. Sharma** (<https://loop.frontiersin.org/people/2203/overview>), Uppsala University, Sweden  
Footnotes

**Reviewed by:**

**Alexander A. Mongin** (<https://loop.frontiersin.org/people/175505/overview>), Albany Medical College, United States

**Ayman ElAli** (<https://loop.frontiersin.org/people/76450/overview>), CHU de Québec Research Center, Canada

**Copyright** © 2018 Zagrean, Hermann, Opris, Zagrean and Popa-Wagner. This is an open-access article distributed under the terms of the **Creative Commons Attribution License (CC BY)** (<http://creativecommons.org/licenses/by/4.0/>). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

**\*Correspondence:** Ana-Maria Zagrean, [ana-maria.zagrean@umfcd.ro](mailto:ana-maria.zagrean@umfcd.ro) (<mailto:ana-maria.zagrean@umfcd.ro>) Aurel Popa-Wagner, [aurel.popa-wagner@geriatricshealthyaging.com](mailto:aurel.popa-wagner@geriatricshealthyaging.com) (<mailto:aurel.popa-wagner@geriatricshealthyaging.com>)

<sup>†</sup>These authors have contributed equally to this work

Write a comment...

[Add](#)

## PEOPLE ALSO LOOKED AT

### Oscillatory Cortical Activity in an Animal Model of Dystonia Caused by Cerebellar Dysfunction (/articles/10.3389/fncel.2018.00390/full)

Elena Laura Georgescu (<https://loop.frontiersin.org/people/631534/overview>), Ioana Antoaneta Georgescu (<https://loop.frontiersin.org/people/630399/overview>), Carmen Denise Mihaela Zahiu (<https://loop.frontiersin.org/people/595950/overview>), Alexandru Răzvan Șteopăoiă (<https://loop.frontiersin.org/people/590964/overview>), Vlad Petru Morozan (<https://loop.frontiersin.org/people/603459/overview>), Adrian Ștefan Pană (<https://loop.frontiersin.org/people/631335/overview>), Ana-Maria Zăgrean (<https://loop.frontiersin.org/people/165431/overview>) and Daniela Popa (<https://loop.frontiersin.org/people/149732/overview>)

### Chemogenetic Recruitment of Specific Interneurons Suppresses Seizure Activity (/articles/10.3389/fncel.2018.00293/full)

Alexandru Călin (<https://loop.frontiersin.org/people/589483/overview>), Mihai Stancu (<https://loop.frontiersin.org/people/589865/overview>), Ana-Maria Zagrean (<https://loop.frontiersin.org/people/165431/overview>), John G. R. Jefferys (<https://loop.frontiersin.org/people/27586/overview>), Andrei S. Ilie (<https://loop.frontiersin.org/people/598163/overview>) and Colin J. Akerman (<https://loop.frontiersin.org/people/11315/overview>)

### Human Brain/Cloud Interface (/articles/10.3389/fnins.2019.00112/full)

Nuno R. B. Martins (<https://loop.frontiersin.org/people/610289/overview>), Amara Angelica (<https://loop.frontiersin.org/people/51997/overview>), Krishnan Chakravarthy (<https://loop.frontiersin.org/people/608131/overview>), Yury Svidinenko, Frank J. Boehm (<https://loop.frontiersin.org/people/612669/overview>), Ioan Opris (<https://loop.frontiersin.org/people/25451/overview>), Mikhail A. Lebedev (<https://loop.frontiersin.org/people/3821/overview>), Melanie Swan (<https://loop.frontiersin.org/people/615699/overview>), Steven A. Garan (<https://loop.frontiersin.org/people/610340/overview>), Jeffrey V. Rosenfeld (<https://loop.frontiersin.org/people/193233/overview>), Tad Hogg (<https://loop.frontiersin.org/people/691054/overview>) and Robert A. Freitas (<https://loop.frontiersin.org/people/11283/overview>)  
Abstract

### Automatic Human Sleep Stage Scoring Using Deep Neural Networks (/articles/10.3389/fnins.2018.00781/full)

Alexander Malafeev (<https://loop.frontiersin.org/people/606445/overview>), Dmitry Laptev (<https://loop.frontiersin.org/people/256618/overview>), Stefan Bauer (<https://loop.frontiersin.org/people/602237/overview>), Ximena Omlin, Aleksandra Wierzbicka (<https://loop.frontiersin.org/people/618344/overview>), Adam Wichniak, Małgorzata Jernajczyk, Robert Riener, Joachim Buhmann (<https://loop.frontiersin.org/people/288404/overview>) and Peter Achermann (<https://loop.frontiersin.org/people/11283/overview>)  
Communication

## Tool

### Editorial: Augmentation of Brain Function: Facts, Fiction and Controversy (/articles/10.3389/fnsys.2018.00045/full)

Mikhail A. Lebedev (<https://loop.frontiersin.org/people/3821/overview>), Ioan Opris (<https://loop.frontiersin.org/people/25451/overview>) and Manuel F. Casanova (<https://loop.frontiersin.org/people/3604/overview>)  
Remodeling In

Response to Stroke

Neurovascular Unit-  
Derived Exosomes  
in Response to  
Stroke

The Potential Use of  
Mesenchymal Stem  
Cells and Their  
Exosomes for Stroke  
Therapy

Conclusion and  
Perspectives

Author  
Contributions

Funding

Conflict of Interest  
Statement

Footnotes

References

(<https://spotlight.frontiersin.org/how-to-enter/>)

utm\_source=FWEB&utm\_medium=FMAIN&utm\_campaign=SPOTLIGHT&utm\_term=BANNER&utm\_content=WEB

Want to win \$100.000 to host your own conference? (<https://spotlight.frontiersin.org/how-to-enter/>)

utm\_source=FWEB&utm\_medium=FMAIN&utm\_campaign=SPOTLIGHT&utm\_term=BANNER&utm\_content=WE

We use cookies on this website to improve your user experience. [Learn more](https://www.frontiersin.org/legal/list-of-cookies) | [Privacy Policy](https://www.frontiersin.org/legal/privacy-policy) | [Close](#)

[Suggest a Research Topic](https://spotlight.frontiersin.org/how-to-enter/?utm_source=FWEB&utm_medium=FMMAIN&utm_campaign=SPOTLIGHT&utm_term=BANNER&utm_content=WEB) ([https://spotlight.frontiersin.org/how-to-enter/?utm\\_source=FWEB&utm\\_medium=FMMAIN&utm\\_campaign=SPOTLIGHT&utm\\_term=BANNER&utm\\_content=WEB](https://spotlight.frontiersin.org/how-to-enter/?utm_source=FWEB&utm_medium=FMMAIN&utm_campaign=SPOTLIGHT&utm_term=BANNER&utm_content=WEB))

[Home](https://www.frontiersin.org/Journal/Frontiers.aspx) (<https://www.frontiersin.org/Journal/Frontiers.aspx>)  
[About Frontiers](https://www.frontiersin.org/about) (<https://www.frontiersin.org/about>)  
[Journals A-Z](https://www.frontiersin.org/about/journalseries) (<https://www.frontiersin.org/about/journalseries>)  
[Institutional Membership](https://www.frontiersin.org/about/Institutional_Membership) ([https://www.frontiersin.org/about/Institutional\\_Membership](https://www.frontiersin.org/about/Institutional_Membership))  
[Contact](https://www.frontiersin.org/about/contact) (<https://www.frontiersin.org/about/contact>)  
[News](https://blog.frontiersin.org) (<https://blog.frontiersin.org>)  
[Submit](https://www.frontiersin.org/submitinfo) (<https://www.frontiersin.org/submitinfo>)  
[FAQs](https://frontiers.zendesk.com/hc/en-us) (<https://frontiers.zendesk.com/hc/en-us>)  
[Terms & Conditions](https://www.frontiersin.org/TermsandConditions.aspx) (<https://www.frontiersin.org/TermsandConditions.aspx>)  
[Privacy Policy](https://www.frontiersin.org/legal/privacy-policy) (<https://www.frontiersin.org/legal/privacy-policy>)  
[Newsletters](http://connect.frontiersin.org/subscriptions/subscribe) (<http://connect.frontiersin.org/subscriptions/subscribe>)  
[RSS/Twitter](https://blog.frontiersin.org/2013/11/01/frontiers-social-media-and-rss/) (<https://blog.frontiersin.org/2013/11/01/frontiers-social-media-and-rss/>)  
[Careers](https://www.frontiersin.org/Careers) (<https://www.frontiersin.org/Careers>)

## TABLE OF CONTENTS

Abstract

© 2007 - 2019 Frontiers Media S.A. All Rights Reserved

Introduction

Exosomes' as a  
Biological  
Communication  
Tool

Neurovascular Unit  
Remodeling in  
Response to Stroke

Neurovascular Unit-  
Derived Exosomes  
in Response to  
Stroke

The Potential Use of  
Mesenchymal Stem  
Cells and Their  
Exosomes for Stroke  
Therapy

Conclusion and  
Perspectives

Author  
Contributions

Funding

Conflict of Interest  
Statement

Footnotes

References